

日本のミナミマグロ延縄操業パターンの変化の検証：2016年漁期

Change in operation pattern of Japanese southern bluefin tuna
longliners in the 2016 fishing season.

伊藤智幸

Tomoyuki ITOH

国立研究開発法人水産研究・教育機構 国際水産資源研究所

National Research Institute of Far Seas Fisheries,

Japan Fisheries Research and Education Agency

要約

日本延縄船によるミナミマグロの漁獲データは、CCSBTにおけるミナミマグロの資源評価ならびにMPにおいて最も重要なものとして使用される。過去10年間と比較し、最近年の操業パターンの変化を検討した。漁獲量、隻数、操業のあった時空間、海区別割合、体長組成、操業の集中度を検討した結果、操業パターンの大きな変化は2016年には生じていなかった。2016年の日本延縄漁業のCPUEは従来と同程度に資源を反映したものとみなすことができるだろう。2016年の漁獲枠増加は、CPUEの増加に貢献し、操業時空間の拡大及び操業回数の増加にはそれより小さい程度で貢献した。

Summary

The Japanese longline data have been used as the most important scientific data for the stock assessment and Management Procedure of southern bluefin tuna (SBT) in CCSBT. Operation pattern of the longline fishing was examined by comparison between the most recent year and previous 10 years. No remarkable change was found in the 2016 operational pattern in terms of catch amount, the number of vessels, time and area operated, proportion by area, length frequency, and concentration of operations. It can be said that the Japanese longline CPUE in 2016 represents the change of SBT stock abundance in consistently as in previous years. The increase of Japanese total catch contributed largely on higher CPUE and contributed lesser degree for the expansion of time and area of operation or increase in the number of operation.

Introduction

CCSBT におけるミナミマグロ *Thunnus maccoyii* の資源評価は漁業情報に大きく依存している。漁業データは、漁業独立調査よりも比較的容易に低コストで広い時空間の情報を集めることができる長所がある。しかしそのデザインが系統だっではないことから、漁業研究者はデータを適切に解釈し、資源指標の変化の理由について資源の変化なのかそれとも漁業の操業パターンの変化に起因するバイアスなのかを区別する必要がある。毎年の注意深い漁業データのモニタリングは CCSBT をはじめとしたまぐろ RFMO での資源評価に不可欠な作業である。

CCSBT におけるそうした操業パターン変化の検証は、ミナミマグロに関連する全ての漁業について毎年実施すべきである。中でも日本延縄船によるミナミマグロの漁獲データは、CCSBT におけるミナミマグロの資源評価において最も重要なものとして使用されてきたこと、ならびに 2011 年に運用を開始した管理方策 MP では、TAC を決定するインプットデータであることから、注意深いモニタリングが必要である。よって我々は毎年、検証文書を提出してきた (e.g. Itoh 2015, Itoh and Yamasaki 2016)。本文書は、2016 年末までのアップデートであり、過去 10 年間との比較で 2016 年の変化を検討する。

The stock assessment of southern bluefin tuna *Thunnus maccoyii* (SBT) in CCSBT relies on data from various fisheries. Fishery data have a crucial advantage that a large amount of information from a large scale in time and space can be easily collected without expensive cost compare to fishery independent researches. However, because its design is not systematic, fishery scientists need to interpret the data properly and to distinguish the reason of any changes in stock indices is whether by changes in stock itself or by bias caused by any change of fishing operational pattern. Careful monitoring of fishery every year is an essential task for stock assessment and management of fish in tuna RFMOs, including CCSBT.

Such a careful examination of fishing operational pattern should be carried out every year for each of the fishing fleets whose data can be used for the stock assessment of SBT in CCSBT. Particularly, it is important for catch data of Japanese longliners, because it has been used as the most important data for the stock assessment of SBT for a long period of years and it is a major input datasets of the Management Procedure which implemented CCSBT in 2011. We have been evaluating operational pattern of Japanese longline for SBT every year (e.g. Itoh 2015, Itoh and Yamasaki 2016). This paper is the update of it up to the end of 2016.

Material and method

二つのデータセットを解析した。一つ目のデータセットは 2001 年から 2016 年までの RTMP データである。Logbook が日本の漁獲についての公式統計ではあるが、Logbook データにはミナミマグロを対象としない操業も含まれ、またデータが使用可能となるまでに 2 年程度の遅れがあることから、最近年の RTMP データと同じ条件で比較することができない。なお、RTMP はミナミ

マグロを対象として出漁する日本延縄船の全てが報告しており、日本が CCSBT 統計海区の 4-9 海区において漁獲するミナマグロ漁獲尾数全体の過去においては 90%以上 (2001-2005 年の平均値は 93%)、近年は 100%が RTMP に含まれていることから、日本延縄船のミナマグロ操業を十分に反映している(Sakai et al. CCSBT/0909/SBT Fisheries-Japan)。2016 年以前の 10 年間 (2006 年-2015 年) を比較対象とした。ある月の緯度 5 度、経度 5 度区画をセルと称し、解析の単位とした。

別データセット (日本タイプ延縄操業別データと称す) も解析に用いた。このデータセットはコア船 CPUE を求めるために作成した、コア船を抽出する前のもので、日本の logbook データを含み (最近年は RTMP 主体)、NZ ジョイントベンチャーや豪州枠の RTMP 船操業も含む shot-by-shot データから構成される (Itoh and Takahashi 2016)。

日本タイプ延縄操業別データを使用してセル数を、全操業並びに SBT (4 歳以上魚) が漁獲された操業について集計した。また同じデータセットを使用して、5 度区画・月内において操業のあった 1 度区画の数も集計した。これは一種の操業海域の集中を示す情報である。

操業海域の集中の度合いは別途開発したインデックスでも示した (Attachment 1)。この集中指数は Dr. Hillary が計算方法を提供したもので、操業緯経度の平均位置に対する分散を指標化したものである。集中指数は日本タイプ延縄操業別データセットに適用し、4-9 海区、4 月-9 月のデータにおける釣鈎数とミナマグロ尾数 (4 歳以上魚) の両方について求めた。

Two datasets were used for analyses in this paper. One dataset used was the RTMP (RealTime Monitoring Program) data between 2001 and 2016. While logbook data are the Japanese official catch-and-effort data, logbook data, which includes longline operations targeting for other tuna species and requires about two years to be available, does not allow comparison with the most recent year to previous years in same condition. RTMP data fully represents Japanese SBT longline operation because it includes all of longline operations targeting for SBT. The data includes more than 90% of total SBT catch of Japan in CCSBT statistical area (Area) 4-9 in the past (mean of 2001-2005 is 93%) (Sakai et al. CCSBT/0909/FisheriesJapan), and 100% of the catch in recent years. Various statistics of the data in 2016 were compared to previous 10 years, 2006-2015. Five degrees longitude, five degrees latitude in a month is defined as one “cell”.

Another dataset (“Japanese-type longline shot-by-shot dataset”) was also used. This dataset was made for the CPUE of core vessels which comprised of Japanese logbook data (mostly RTMP data in the most recent year), RTMP data from Australia in the 1990s, and New Zealand charter vessel data (Itoh and Takahashi 2017). Data of all vessels, before extract the core vessels, were used for analysis.

The numbers of cells were calculated for the Japanese-type longline shot-by-shot dataset in two cases, all operations and operations with catch of SBT age 4+. In addition, the number of one degree square operated within five degrees square, a kind of indicator for

operational concentration, was calculated using this dataset.

Concentration of operation was also calculated as another index (Attachment 1). The concentration index was derived from variance to the mean latitude and longitude of operations over the years, and the method was provided by Dr. Hillary in CSIRO. The index was calculated in two cases, all operations and operations with catch of SBT age 4+ by using the Japanese-type longline shot-by-shot dataset in Areas 4 to 9 in April-September.

Result

1. 2016年の漁獲、努力量、サイズの概要 Summary of the catch, effort and size in 2016

図1に、CCSBT統計海区4-9海区における、毎年の隻数、使用鈎数、ミナミマグロ漁獲尾数を、2006年から2015年までの平均値に対する相対値で示す。どの値も2005年から2008年または2010年まで減少し、2010年から2013年までは横ばいで推移した。その後2016年に掛けて、ミナミマグロ漁獲尾数は大きく増加（2013年の1.63倍）し、使用鈎数はわずかに増加（2013年の1.40倍）し、隻数はほぼ横ばいである。

図2に、隻数、使用鈎数、ミナミマグロ漁獲尾数の統計海区別組成を示す。漁獲尾数では、7海区の割合が2004年に大きく減少し、2006年から2011年に掛けて再び増加した。それに伴って4海区と8海区の割合が減少した。2012年以降の年別の海区組成は類似している。

図3にミナミマグロの体長組成を示す。2016年の体長組成は130cmFLと145cmFLに二つのピークを持っていた。2015年には125cmFLと145cmFLにピークを持っており、全体的に類似しているが、わずかに大型化した。小型魚の部分は放流による影響も受けている点に留意すべき(Itoh et al. 2014 CCSBT-OMMP/1406/08)。

Figure 1 shows relative values of the numbers of vessels, hooks used and SBT caught to the mean values between 2006 and 2015 in the Areas 4 to 9. The values decreased from 2005 to 2008 or 2010 followed by stable period up to 2013. Up to 2016, the number of SBT caught largely increased (1.63 times of 2013), the number of hooks used slightly increased (1.40 times of 2013), and the number of vessels was stable.

Figure 2 shows the compositions of numbers of vessels, hooks used and SBT caught by Area. The proportion of number of SBT in Area 7 were dropped in 2004 and increased again from 2006 to 2011. Along with the increase of proportion in Area 7, those in Area 4 and Area 8 were decreased. Since 2012, composition by Area has been similar over years.

Figure 3 shows fork length frequency of SBT by year. In 2016, it has two peaks in 130 cmFL and 145 cmFL. In 2015, it had two peaks in 125 cmFL and 145 cmFL. Length frequency in 2016 was similar to that in 2015, but slightly shifted toward larger. Note that frequencies in small fish are affected from small fish release and discards (Itoh et al. 2014 CCSBT-

2. 操業時空間の変化 Changes of the time and space operated

図 4 に、RTMP 船データセットにおける 4-9 海区内の操業のあったセル (5x5 度・月単位) 数の変化を示す。合計セル数は 2006 年の 165 セルから 2014 年の 94 セルまで減少し、2015 年には 125 セルに増加したが 2016 年には 97 セルに低下した。

表 1 に、セル数を年、月、海區別に示す。表 2 には、その操業回数を示す。過去 5 年間または 10 年間と比較して 2016 年の操業時期、海域はほぼ同様であった。詳細に見ると、全海区・一年間に対する海区・月別の操業割合が過去 5 年間の組成に対して 3%以上変動したのは、7 海区 5 月 (増加)、9 海区 5 月 (増加) であった。各海区の中心時期は 4 海区 5 月、7 海区 4 月、8 海区 8 月、9 海区 6 月であり、過去と同様であった。8 海区では 2010 年以降、4 月から 5 月の操業割合が増加しているが、主漁期である 8 月よりは依然として低い。

図 5 に RTMP 船データセットにおける 1 セル当たりの操業回数を示す。日本の総漁獲枠が半分以下に減少し、漁期撤廃に伴ってセル数が増加したことの結果として、1 セル当たりの操業回数は 2005 年の 114.3 回/セルから 2010 年に 20.7 回/セルにまで減少した。その後わずかな増加が継続し、2016 年には 47.7 回/セルとなった。7 海区では 1 セル当たりの操業回数が 2010 年から大きく増加し、2013 年からは変動している。

図 6 は日本タイプ延縄操業別データセットを使って求めた 4-9 月、4-9 海区内の操業セル数である。RTMP に基づく図 4 よりも長期データであるが、RTMP データが主体となる最近年 (2016 年) とそれ以前との一貫性には注意が必要である。SBT が漁獲されなかった操業も含めた全操業で見ると、操業セル数は 1980 年代から次第に減少し、2002 年、2003 年に低くなった後に 2006 年にかけて増加したが、その後はまた減少し、2013 年以降は横ばいである (Fig. 6a 上図)。セルの定義を 5 度区画・月から 1 度区画・月に変更しても傾向は変わらない (Fig. 6a 中図)。5 度区画・月の 1 セル当たりの操業回数 (Fig. 6a の白丸) は 1986 年から 2005 年まではほぼ一定であったが、その後は減少し、2007 年以降は横ばいである。2010 年以降はわずかに増加傾向である。セル数の減少は特定の海区だけでなく、全ての海区で生じていた。

データをミナミマグロ 4 歳以上魚が漁獲された操業に限定しても、全操業で見られたものと傾向に違いはなかった (Fig. 6b)。

Figure 4 shows the change of the number of cells (five degree square and month) in Area 4-9 in the RTMP data. The total number of cells has been decreasing from 165 in 2006 to 94 in 2014. It turned increasing in 2015 as 125 cells, however, decreased again to 97 in 2016.

Table 1 shows the number of cells by year, month and Area. Table 2 shows the number of operations. Fishing season and area in 2016 were similar to those in previous five or 10 years in general. Seeing in detail, it increased (>3% of previous five years composition) in May in Area 7 and May in Area 9 in terms of the proportions in the number of

operations by Area and month to the total number of operations in Area 4-9 and Month 4-9 in 2016. The main fishing periods were same as previous years like May in Area4, April in Area 7, August in Area 8, and June in Area 9. While the proportion in the number of operation were higher from April to May than in previous years, it was lower than that in August, the main fishing season, in Area 8.

Figure 5 shows the number of operations per cell in the RTMP dataset. Because the allocation of TAC to Japan was reduced to less than half and the number of cell operated was increased with the lift of seasonal area closure, the number of operations per cell has decreased to 20.7 times per cell in 2010 from 114.3 times per cell in 2005. Since then, slight increase has continued, and it reached 47.7 times per cell in 2016. In Area 7, the number of operations per cell increased to 2010 and fluctuated since then.

Figure 6 shows the number of cells operated in Area 4-9 and month 4-9 in the Japanese-type longline shot-by-shot dataset. While the time series was longer than the RTMP dataset, it should be noted that most recent year (2016) was based on RTMP and may have inconsistency to previous years. In all operations including SBT zero catch, the number of cells decreased since the 1980s to 2002 and 2003 followed by slight increase until 2006, decreasing further, and then stable since 2013 (Fig. 6a upper panel). No difference was found in the case that the cell was defined as one degree square and month (Fig. 6a middle panel). The number of operations per cell in five degree (line with open circle in Fig. 6a) had been stable between 1986 and 2005, then decreased and has been stable since 2007, but has been slightly increasing since 2010. Decrease of the number of cells was observed in all Areas.

No difference was found when the data was limited in operations with catch of SBT 4+ only, instead of all operations (Fig. 6b).

3. 操業の集中度 Concentration of area operated

日本タイプ延縄操業別データによる、ある5度区画内で操業のあった1度区画の数(全数は25)は、1986-2006年の平均値は6.9個、2007-2012年には低下してほぼ横ばいとなり、2013年から2016年には増加した (Fig. 6a 下図)。データをミナミマグロ4歳以上魚が漁獲された操業に限定した場合、2007年に低下してから横ばいのままである (Fig. 6b 下図)。

集中度指数は海域別に求めた (図7)。値が高いことは操業海域の拡散を、また低いことは操業海域の集中を意味する。Area 6は2016年にデータが無かった。経年的には8海区、9海区は安定的に推移し、5海区、7海区は変動が大きかった。8海区では2013年と2014年に、漁獲尾数では集中しているが努力量の分布は拡散していた。操業位置を詳細に検討したところ、7海区は、2003年まではタスマニア島の東西両方で操業があったが、2004年以降は東側でしか操業してお

らず、全年からもとめた 7 海区の漁場の中心から離れたデータが多いことによって指数が増加 (拡散) した。

The number of one degree square operated in a five degrees square (total is 25) was 6.9 in the average of 1986-2005. It dropped and was stable in 2007-2012, then increasing from 2013 to 2016 (Fig. 6a bottom panel). When the data was limited in operations with SBT 4+ only, it dropped in 2007 and stable up to 2016 (Fig. 6b bottom panel).

Figure 7 shows the concentration index by Area. Smaller and larger values indicate more and less concentrated operation respectively. No data were available for Area 6 in 2016. The time series have been stable in Area 8 and Area 9, and fluctuated largely in Area 5 and Area 7. In Area 8 in 2013 and 2014, it was less concentrated in terms of hooks while it was still concentrated in catch. Detailed examination of operated area found that, in Area 7, there were operations both the east and west of Tasmania Island before 2003, but only in the east since 2004 (there were only data far apart from the center of Area 7 which obtained from all the years) and resulted in increase of the index (less concentrate).

4. 船の一貫性 Vessel consistency

表 3 に、2016 年の RTMP 参加船 (かつミナミマグロを漁獲した船) が過去とどれほど共通しているかを示す。2016 年の 88 隻中、49 隻は 2001-2005 年に RTMP においてミナミマグロ操業を実施したことがあり、2001-2005 年に 4 年間または 5 年間実施した船が 56% と依然として大きな割合を占めており、船は一貫したものが多いことが分かる。

Table 3 shows the consistency of the vessels that participated the RTMP in 2016 (and caught any SBT) with those in 2001-2005. Among 88 vessels in 2016, 49 vessels caught SBT in 2001-2005 RTMP and they still consist of a large part of SBT vessels (56%).

Discussion

漁獲量、隻数、操業のあった時空間、海區別割合、体長組成、操業の集中度を検討した結果、操業パターンの大きな変化は 2016 年には生じていなかった。2016 年の日本延縄漁業の CPUE は従来と同程度に資源を反映したものとみなすことができるだろう。

7 海区での操業割合は、10 年間継続して増加している。低水準であった 2000 年級が加入したころ、タスマン海ではミナミマグロの漁況が悪く、日本延縄船が 7 海区での操業を避けていた。新たな増加というよりはそれらの漁船団が戻ったものである (Itoh et al 2015 CCSBT-ESC/1509/SBT Fisheries -Japan)。

日本の漁獲量は 2013 年の 2694 トンから 2014 年の 3371 トン、2015 年に 4745 トン、2016 年に 4721 トン (2013 年の 1.75 倍) へと増加した。これに対して RTMP データにおける 2016 年の 4-9 海区の操業回数は 2013 年に対して 1.40 倍にしか増えなかった。そして操業セル数は 0.97 倍 (97/100) にしか増えなかった。漁獲量の増加の多くは CPUE の増加 (1.63 倍) が貢献してお

り、操業回数や操業時空間の拡大への貢献はそれより小さかった。

No remarkable change was found in the 2016 operational pattern in terms of catch amount, the number of vessels, time and area operated, proportion by area, length frequency, and concentration of operations. It can be said that the Japanese longline CPUE in 2016 represents the change of SBT stock abundance in consistently as in previous years.

Proportion of operation in Area 7 has been increasing in last 10 years. In the early 2000s, many Japanese longline fleets escaped from Area 7 with its low CPUE probably due to the weak cohort around the 2000 year classes (Itoh et al. 2015 CCSBT-ESC/1509/SBT Fisheries-Japan).

The total catches of Japan were increased from 2694 tons in 2013 and 3371 tons in 2014, to 4745 tons in 2015 and 4721 tons in 2016 (1.75 times of 2013). However, the factor in the number of operations in 2016 was small as 1.40 times of 2013 in RTMP data. The factor in the number of cells operated was also small as 0.97 times (97/100) of 2013. The increase of total catch contributed largely on higher CPUE (1.63 times of 2013) and contributed lesser degree for the expansion of time and area of operation or increase in the number of operation.

References

- Itoh T. 2015 Change in operation pattern of Japanese southern bluefin tuna longliners in 2014. CCSBT-ESC/1509/30.
- Itoh, T., K. Suzuki and O. Sakai. 2014 Mortality estimation for southern bluefin tuna released and discarded from Japanese longline fishery. CCSBT-OMMP5/1406/08.
- Itoh, T. and N. Takahashi 2016. Update of the core vessel data and CPUE for southern bluefin tuna in 2016. CCSBT-CPUE/1606/04.
- Itoh T. and Izumi Yamasaki 2016 Change in operation pattern of Japanese southern bluefin tuna longliners in the 2016 fishing season. CCSBT-ESC/1609/22.
- Itoh, T., Izumi Yamasaki and Sayako Takeda. 2015. Review of Japanese southern bluefin tuna fisheries in 2014. CCSBT-ESC/1509/SBT Fisheries-Japan.
- Sakai, O., T Itoh and T. Sakamoto. 2009. Review of Japanese SBT Fisheries in 2008. CCSBT-ESC/0909/Fisheries-Japan.

Table 1. Number of 5x5 degree square where longline operations conducted by year, month and area in RTMP data.

Area	Month	Year													
		2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
4	2								1			1			
	3							3	2	1	1	3	2	1	
	4	1	2	2		2	4	2	2	2	3	1	2		2
	5	3	2	2	2	3	2	5	3	4	4	2	3	2	2
	6	4	5	5	3	5	5	5	6	6	3	4	3	2	4
	7	4	6	6	4	6	5	2		2		2	1	3	1
	8		3	1	1	1	1	3	2	2		2	2	1	
	9							1	1	1	1				
	10	4					1	2			1			1	
	11	4													
	12	4												1	
	5	7		1	1	2	2	2	2	2	1	1	2		
8					3	3	2	2		2		2		2	2
9						2	2			1				2	
10							1			1					
12											2				
6	4					2									
	5					2									
	6				1	1									
7	1						2								
	2								2						
	3								1	2	2	2	2	2	2
	4	6	2	2		2	2	2	2	2	3	2	2	2	2
	5	2	2	2	3	2	2	2	2	2	2	2	2	2	2
	6	2	2	2	2	2							2		2
	7	1			1										
	9	2			2										
	10	1	1			1									
	11		1			1									
	8	1					1	2	2	6		1			
3							2					1	1	4	3
4							2		3	1	1	4	4	6	6
5		1	3	5	1		3		7	4		5	3	4	3
6			4	5					2	2	3	2	3	2	2
7					5	8	6	4	2	1	5	3	2	3	2
8					8	8	6	6	4	5	7	5	3	4	3
9		11	13	5	7	6	5	4	2	3	2	3	2	1	3
10		13	9	7	8	8	8	8	2	6	2		2		
11		9	9	8	8	7	7	6	3	4	1	1	2	4	
12		7	8	4	7	6	4	6	1	1			2	4	
9		2								1					
	3							3	7	5	5	2	3	3	
	4						5	7	8	15	8	13	6	8	8
	5	14	19	25	21	16	14	8	20	15	13	5	10	16	13
	6	17	23	20	18	18	15	10	14	13	12	12	11	15	8
	7	15	19	19	21	16	16	12	12	11	9	8	10	11	8
	8		12	13	15	8	11	9	9	9	6	5	6	4	7
	9				11	5	5	3	5	6	4	6	3	7	7
	10				7	4	6	1	3	2	3	4	1	6	4
	11				4	2	2		2		1	2			
	12						1	1	1						

Dotted line shows 2006 when the individual quarter system started.

Table 2. Number of operations by year, month and area in RTMP data

Area	Month	Year										%10Y	%5Y	2016%					
		2003	2004	2005	2006	2007	2008	2009	2010	2011	2012				2013	2014	2015	2016	
4	2								4				8				0.0%	0.0%	0.0%
	3							30	33	3	2	31	5	1			0.2%	0.2%	0.0%
	4		2	23	13		39	45	47	60	30	34	2	38		10	0.6%	0.6%	0.2%
	5		347	447	731	530	55	140	203	139	131	149	38	64	108	44	3.2%	2.8%	1.0%
	6		1015	1179	1122	457	324	147	46	52	74	8	55	52	37	81	2.6%	1.3%	1.7%
	7		911	1110	732	115	75	49	35		72		39	14	46	10	0.9%	1.0%	0.2%
	8			10	1	3	2	1	3	15	5		34	16	2		0.2%	0.3%	0.0%
	9								12	4	4	35	4	3			0.1%	0.2%	0.0%
	10		96				2	6			2				1		0.0%	0.0%	0.0%
	11		120														0.0%	0.0%	0.0%
	12		47														0.0%	0.0%	0.0%
	5	7			2	6	11	22	17	8	25	4	1	10		3	0.2%	0.1%	0.1%
8						27	34	23	11		65		10		30	14	0.4%	0.6%	0.3%
9							17	7			20				6		0.1%	0.1%	0.0%
10								2			3						0.0%	0.0%	0.0%
6	4																0.0%	0.0%	0.0%
	5																0.1%	0.0%	0.0%
	6					1	22										0.0%	0.0%	0.0%
7	1							2									0.0%	0.0%	0.0%
	2																0.0%	0.0%	0.0%
	3									7							0.0%	0.0%	0.0%
	4		648	530	603		27	181	189	290	449	547	522	500	613	574	6.9%	14.8%	12.4%
	5		1032	646	397	180	352	183	64	61	118	48	399	276	243	457	4.0%	6.1%	9.9%
	6		254	2	3	66	92							14		35	0.4%	0.1%	0.8%
	7		1			46											0.1%	0.0%	0.0%
	9		47			6											0.0%	0.0%	0.0%
	10		1	6			1										0.0%	0.0%	0.0%
	11			13			2										0.0%	0.0%	0.0%
	8	1						1	2	4	60		2					0.1%	0.0%
3								30					7	6	25	23	0.1%	0.2%	0.5%
4								5		34	3	16	129	206	138	171	1.1%	2.8%	3.7%
5			1	411	651	12		26		55	26		180	246	214	157	1.6%	3.8%	3.4%
6				104	13					88	16	33	17	154	85	33	0.8%	1.7%	0.7%
7						76	103	177	33	4	15	55	47	17	116	30	1.3%	1.4%	0.6%
8						407	773	921	988	259	197	425	461	600	716	738	12.0%	13.5%	15.9%
9			961	489	551	270	630	482	251	13	214	156	76	58	144	265	4.8%	3.7%	5.7%
10			842	589	687	343	379	131	163	8	68	7		7			2.3%	0.5%	0.0%
11			825	904	821	518	750	362	357	95	125	16	13	32	28		4.8%	1.2%	0.0%
12			316	618	488	259	115	177	280	2	39			17	25		1.9%	0.5%	0.0%
9		2									1					9		0.0%	0.1%
	3							23		68	44	66	19	12	10		0.5%	0.9%	0.0%
	4						66	111	46	298	155	196	107	113	141	138	2.6%	4.0%	3.0%
	5		2564	2383	1897	905	160	220	184	473	459	359	241	236	358	573	7.5%	9.3%	12.4%
	6		2672	2826	2537	1163	575	792	394	443	468	534	504	476	655	640	12.5%	14.9%	13.8%
	7		586	2832	2802	1513	683	1032	436	222	270	351	179	349	199	444	10.9%	7.6%	9.6%
	8			522	1261	1738	755	623	266	115	86	116	31	25	64	37	7.9%	1.8%	0.8%
	9					1033	623	224	34	22	121	38	33	55	80	39	4.7%	1.8%	0.8%
	10					262	183	105	50	22	26	16	55	1	31	27	1.6%	0.7%	0.6%
	11					24	5	30		35		7	11				0.2%	0.1%	0.0%
	12							5	1	1							0.0%	0.0%	0.0%
	Total		13288	15646	15316	9965	6911	6323	4097	3041	3368	3298	3275	3637	4161	4631	100.0%	100.0%	100.0%

“%10Y” and “%5Y” are the proportion of the Area and month to the sum of 10 years (2006-2015) and 5 years (2011-2015), respectively. “2016%” is the proportion of the Area and month to the sum of 2016 data. Shadow and line enclosing denotes 2016% is higher (>3%) than %5Y.

Table 3. Number of vessels that caught SBT in RTMP between 2006 and 2016 by the number of years participated in RTMP in past years (2001-2005) in RTMP data.

	Number of years participate in the RTMP during 2001-2005					
	0 year	1 year	2 year	3 year	4 year	5 year
2006	5	7	15	9	20	67
2007	5	10	16	10	22	74
2008	11	8	16	8	22	61
2009	13	4	13	6	15	49
2010	9	4	12	5	14	42
2011	8	6	12	6	12	39
2012	11	6	13	6	14	43
2013	13	4	11	4	14	42
2014	18	3	10	6	13	40
2015	20	3	10	6	11	39
2016	22	3	8	6	11	38




Fig.1. Changes in the number of operation, the number of vessels and the number of SBT caught in Area 4-9 in 2016 and previous years in RTMP data. Y axis is the relative value to the average of previous 10 years.






Fig.2. Proportions of Area in the number of vessels, the number of hooks used and the number of SBT caught in 2016 and previous years in RTMP data.




Fig.3. Length frequency distributions of SBT by year in 2016 and previous years in RTMP data.




Fig.4. Changes of the number of cells (5 degrees latitude and longitude and month) operated in 10 years in Area 4-9 in RTMP data.




Fig.5. Changes of the number of longline operations per cell (5 degrees latitude and longitude and month) in 10 years in Area 4-9 in RTMP data.




Figure 6a. Number of cells in the Japanese-type longline dataset for all operations.
 (Top panel) Bar represents the number of 5x5 degrees square and month (cell) where fishing operated by CCSBT statistical area and refer to left side y-axis. Line with circle plot represents the mean annual number of operations per cell and refer to right side y-axis. (Middle panel) Bar represents the number of 1x1 degree square and month (cell) where fishing operated by CCSBT statistical area and refer to left side y-axis. Line with circle plot represents the mean annual number of operations per cell and refer to right side y-axis. (Bottom panel) Composition of frequency for the number of 1x1 degree square and month cells operated in a 5x5 degree squares and month cell. Refer to left side y-axis. The grey band is one of 25 cells and that at top is 25 of 25 cells, and every five is colored. Line with triangle represents the mean number of 1x1 month cells operated in a 5x5 month cell and refer to right side y-axis.




Figure 6b. Number of cells in the Japanese-type longline dataset for operations of SBT 4+ catch positive. See explanation in Fig. 6a.




Figure 7. Concentration index in the Japanese-type longline dataset for catch (age 4+) or hooks to years by CCSBT statistical area
Smaller/larger values relate to more/less aggregated.

SBT exploratory data analysis Idea

In conjunction with usual analyses done for the indicators paper there are some potentially revealing but simple things we can do to be primed for the upcoming assessment of SBT.

CPUE data

This obviously really refers to the LL_1 CPUE data that will form the basis of the main abundance index in the OM. While that will work with the standardised CPUE there are several things we can look at in the raw data. In terms of a reasonably detailed spatial analysis, by area and 5×5 square there are some basic spatial statistics we can look at, henceforth, I_{ijk} denotes the raw CPUE (over a given length/age range) for year i , in region j , in square k . To look at a rough trend in how the average fishing location as changed in each area over time we can simply compute the centre of mass, R_{ij} of the CPUE in a given year i and region j :

$$R_{ij} = \frac{\sum_{k \in j} \rho_k I_{ijk}}{\sum_{k \in j} I_{ijk}}, \quad (1)$$

where ρ_k denotes the physical location of square k in area j (i.e. lat and long) and then the square center of mass of that region in a given year is the square containing R_{ij} . A secondary measure is to see how the density of either effort, catch or CPUE changes over time and in each area. This can be calculated by first estimating the discrete mass density of the given quantity of interest, X_{ijk} , in a given region at a given time:

$$\mu_{ijk} = \frac{X_{ijk}}{\sum_{k \in j} X_{ijk}}, \quad (2)$$

and from this we can estimate the (relative) aggregation, α_{ij} , of the quantity X_{ij} fairly easily:

$$\alpha_{ij} = \frac{E^k(\mu_{ijk})^2}{E^k(\mu_{ijk}^2)} \quad (3)$$

where $\alpha_{ij} \in [N_j^{-2}, 1]$ (where N_j is the number of squares in region j) and smaller/larger values of α relate to more/less aggregated spatial quantities. Even spread it is equal to 1 and all in one square it is equal to N_j^{-2} .