



Australia's 2019–2020 Southern Bluefin Tuna Fishing Season

P.I. Hobsbawn and H.M. Patterson

Research by the Australian Bureau of Agricultural and Resource Economics and Sciences

Working Paper CCSBT-ESC/2108/SBT Fisheries—Australia prepared for the CCSBT Extended Scientific Committee for the 26th Meeting of the Scientific Committee, 23 - 31 August 2021

Technical Report 21.6
July 2021



© Commonwealth of Australia 2021

Ownership of intellectual property rights

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

Creative Commons licence

All material in this publication is licensed under a [Creative Commons Attribution 4.0 International Licence](#) except content supplied by third parties, logos and the Commonwealth Coat of Arms.

Inquiries about the licence and any use of this document should be emailed to copyright@awe.gov.au.



Cataloguing data

This publication (and any material sourced from it) should be attributed as: Hobsbawn, PI & Patterson, HM & 2021, *Australia's 2019–20 southern bluefin tuna fishing season*, ABARES, Canberra, August, DOI: [10.25814/1kxa-re78](https://doi.org/10.25814/1kxa-re78).
CC BY 4.0.

ISSN 189-3128

Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES)

GPO Box 858 Canberra ACT 2601

Telephone 1800 900 090

Web awe.gov.au

Disclaimer

The Australian Government acting through the Department of Agriculture, Water and the Environment, represented by the Australian Bureau of Agricultural and Resource Economics and Sciences, has exercised due care and skill in preparing and compiling the information and data in this publication. Notwithstanding, the Department of Agriculture, Water and the Environment, ABARES, its employees and advisers disclaim all liability, including liability for negligence and for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying on any of the information or data in this publication to the maximum extent permitted by law.

Professional independence

The views and analysis presented in ABARES publications, including this one, reflect ABARES professionally independent findings, based on scientific and economic concepts, principles, information and data. These views, analysis and findings may not reflect or be consistent with the views or positions of the Australian Government, or of organisations or groups who have commissioned ABARES reports or analysis. More information on professional independence is provided on the ABARES website at: <https://www.agriculture.gov.au/abares/about/research-and-analysis#professional-independence>

Acknowledgements

The authors thank Matt Daniel (AFMA) and Neil Hughes (DAWE) for their input and comments on this report. Work was funded by Fisheries Resources Research Fund and ABARES.

Contents

Summary	iv
1 Introduction	1
1.1 History	1
1.2 Recent seasons	1
2 Catch and effort	4
2.1 Purse seine fishery	4
2.2 Longline fishery	4
3 Fleet size and distribution	12
3.1 South Australia (SA)	12
3.2 Western Australia (WA)	12
3.3 New South Wales (NSW)	12
3.4 Tasmania (TAS)	12
3.5 Queensland (QLD)	12
4 Research and monitoring to improve estimates of attributable catch	13
4.1 Recreational fishing	13
4.2 Discards in the commercial fishery	15
4.3 Customary and/or traditional fishing	17
4.4 Artisanal fishing	17
5 Other relevant information	18
5.1 Stereo-video commercial trial	18
Appendix A: SBT fishing season dates 1988-89 to 2018-19	19
Appendix B: Purse seine fishing season duration	20
Appendix C: Australian surface catch for farm operations, 1994-95 to 2018-19	21
Appendix D: Nominal CPUE for all Australian longliners, 2018 and 2019 calendar years	22
Appendix E: Summary of observed catch and effort by year and sector	24
Appendix F: Number of biological samples taken in observer programs (year and sector)	28
Appendix G: An overview of the Australian SBT observer program	29
Observer sources and training	29
Purse seine fishery – Great Australian Bight 2018-19	29
Longline fishery	29
Scientific observer program design and coverage	30
Observer data collected	30
Tag return monitoring	32
Appendix H: Australian southern bluefin tuna research projects	33
References	35

Tables

Table 1 Australian Catch (t) by Gear and State for Fishing Seasons 1988–89 to 2019–20.....	2
Table 2 Average fork length (cm) of SBT landed in each Australian state, 1989 to 2020	8
Table 3 Reported tag releases of SBT by Australian recreational fishers, 1990 to 2020.....	15

Figures

Figure 1 Australian SBT catch in the 2019 calendar year. Note that catch is centred in each grid square.	6
Figure 2 Australian SBT catch in the 2020 calendar year. Note that catch is centred in each grid square.	7
Figure 3 Length frequency of SBT purse seine catch in Australian waters raised to total catch, 2012–13 to 2019–20 fishing seasons (source: tow cage size monitoring database).....	9
Figure 4 Length frequency histograms for retained SBT longline catch in Australian waters raised to total catch, 2012–13 to 2019–20 quota years (source: AFMA observer data and processor monitoring data).....	10
Figure 5 Length frequency of SBT measured by Australian observers on Japanese longliners fishing in the Tasmania region, 1988 to 1997. The data include small SBT tagged and released by observers.	11
Figure 6 The number of sets recorded in the SBT purse seine fishery from December 2019 to April 2020. The hatching represents areas where observations took place.....	31

Summary

The 2019–20 Southern Bluefin Tuna (SBT) fishing season report summarises catches and fishing activities in the Australian Southern Bluefin Tuna Fishery up to and including the 2019–20 fishing season¹ (1 December 2019 – 30 November 2020).

Australia's allocation as agreed by the Commission for the Conservation of Southern Bluefin Tuna was 6,165 t for the 2019–20 fishing season. However, this was adjusted to account for overcatch in the previous fishing season so the effective TAC was 6,125 t. A total of 39 commercial fishing vessels landed SBT in Australian waters in the 2019–20 fishing season for a total catch of 5,429 t. A total of 84.1% of the catch was taken by purse seine with the remainder taken by longline, pole-and-line, rod-and-reel and trolling. Seven purse seiners fished off South Australia for the Australian farming operations during the 2019–20 fishing season, with live bait, pontoon-towing and feeding vessels also involved. Most of the purse seine fishing commenced in mid-December 2019 and finished in mid-March 2020.

Length frequency data from the purse seine fishery from 2005–06 to 2006–07 indicated a shift to smaller fish compared to previous years, but this trend has showed signs of reversal since 2007–08, possibly due to the targeting of larger fish. The average length of SBT transferred to farms in South Australia in 2019–20 was 87.8 cm.

In the 2019–20 fishing season, observers monitored 9.9% of purse seine sets where fish were retained for the farm sector and 10.3% of the estimated SBT catch. In 2020, observers also monitored 11.0% of longline hook effort in the Eastern Tuna and Billfish Fishery during the months and in the areas of the SBT migration through that fishery. Observer coverage of longline hook effort in the entire Western Tuna and Billfish Fishery was 12.1% in 2020.

¹ Various time periods, such as 'calendar years', 'fishing seasons' and 'quota years', can be used when describing Australia's SBT Fishery. Unless otherwise indicated, we have used quota years in this report, but note that fishing seasons of the various fishery components often span quota years. The start and end dates of Australian quota years have varied and are presented in Appendix 1.

1 Introduction

This report summarises catches and fishing activities in the 2019–20 fishing season of the Australian Southern Bluefin Tuna (*Thunnus maccoyii*; SBT) Fishery. Caton et al. (1995) provides a more detailed historical description of the fishery.

1.1 History

Troll catches of SBT were reported as early as the 1920s off the east coast of Australia, but significant commercial fishing for SBT commenced in the early 1950s with the establishment of a pole-and-live-bait fishery off New South Wales (NSW), South Australia (SA) and, later (1970) Western Australia (WA). Purse seine gear overtook pole as the main fishing method and catches peaked at 21,500 t in 1982. The bulk of this early Australian catch of SBT was canned. Following quota reductions in 1983–84, the WA pole fishery for very small juveniles closed down and the south-eastern fishery began to target larger juveniles to supply the Japanese sashimi market. Surface catches were further reduced between 1989 and 1995 when about half of the Australian total allowable catch (TAC) was taken by Australia–Japan joint venture longliners in the Australian Fishing Zone (AFZ). The joint ventures ceased in late 1997. From 1992 to 1998, domestic longliners operating off Tasmania (TAS) and NSW also took around 5–10% of the total Australian catch.

In 1990–91, about 20 t of SBT were transferred to fattening cages in Port Lincoln, SA, to enhance their value. Utilisation of the Australian SBT TAC in 'farming' operations increased from 3% of the TAC in 1991–92 to 98% in 1999–2000. The purse-seine fishery now accounts for ~85% of the total catch.

Following declaration of the AFZ in 1979, Japanese longliners fished under a range of bilateral conditions, a real-time monitoring program and joint-venture arrangements until 1997 when Japanese longliners were excluded from all AFZ fishing operations following failure to reach agreement on a global TAC within the Commission for the Conservation of Southern Bluefin Tuna (CCSBT). Caton and Ward (1996) provides copies of annual subsidiary agreements for the operations of bilateral-licensed longliners in the AFZ from 1979–80 to 1994–95.

1.2 Recent seasons

The Australian commercial SBT catches for the 2019 and 2020 calendar years were 6,185 t and 4,757 t, respectively. The catches for the 2018–19 and 2019–20 fishing seasons were 6,074 t and 5,429 t, respectively (Table 1). The TAC for the 2019–20 fishing season was 6,165 t. However, this was adjusted to account for overcatch in the previous fishing season so the effective TAC was 6,125 t. This action is consistent with the guidelines in the CCSBT Resolution on Limited Carry-forward of Unfished Annual Total Available Catch of Southern Bluefin Tuna adopted in October 2017 by the CCSBT. See Appendix 1 for quota year dates and Appendix 2 for duration of the farm sector fishing season.

Australia's 2019–20 Southern Bluefin Tuna Fishing Season

Table 1 Australian Catch (t) by Gear and State for Fishing Seasons 1988–89 to 2019–20

Fishing Season	Western Australia				South Australia				New South Wales			Tasmania			Large Longliners			Australia Total			Total All Gears	
	Albany Pole	Esperance Pole	Long-line	Total	Pole & Purse Seine	Farm Cages	Long-line	Total	Pole & Purse Seine	Long-line	Total	Troll	Long-line	Total	Aust. Charter	Joint-venture	Total	Domestic Surface	Domestic Long-line	Total Long-line		RTMP
1988–89	204	221	0	425	4872	0	0	4872	0	1	1	2	0	2	0	684	684	5299	1	685	0	5984
1989–90	133	97	0	230	4199	0	0	4199	0	6	6	14	0	14	0	400	400	4443	6	406	0	4849
1990–91	175	45	0	220	2588	0	0	2588	0	15	15	57	0	57	255	881	1136	2865	15	1151	300 ^a	4316
1991–92	17	0	0	17	1629	138	14	1781	34	90	124	36	20	56	59	2057	2116	1854	124	2240	800	4894
1992–93	0	0	0	0	716	722	68	1506	16	238	254	23	44	67	0	2735	2735	1477	350	3085	650	5212
1993–94	0	0	0	0	621	1294	55	1970	0	286	286	7	105	112	0	2299	2299	1922	446	2745	270	4937
1994–95	0	0	0	0	908	1954	2	2864	0	157	157	4	109	113	0	1295	1295	2866	268	1563	650	5080
1995–96	0	0	0	0	1447	3362	0	4809	28	89	117	0	262	262	0	0	0	4837	351	351	0	5188
1996–97	0	0	0	0	2000	2498	0	4497	7	229	236	2	242	244	0	0	0	4507	472	472	0	4978
1997–98	0	0	0 ^b	0	916	3488	0 ^b	4403	0 ^c	475	475	0 ^d	219	219	0	0	0	4433	664	664	0	5097
1998–99	0	0	0 ^b	0	28	4991	0 ^b	5018	0 ^c	97	97	0 ^d	116	116	0	0	0	5016	216	216	0	5232
1999–00	0	0	0 ^b	0	0	5130	13	5143	0	114	114	0	0 ^d	0	0	0	0	5130	127	127	0	5257
2000–01	0	0	0 ^b	0	0	5162	6	5168	0	32	32	0	0 ^d	0	0	0	0	5162	38	38	0	5247
2001–02	0	0	7	7	0	5234	0	5234	0	22 ^e	22	0	0 ^d	0	0	0	0	5234	29	29	0	5262
2002–03	0	0	0 ^f	0	0	5375	0	5375	0	17	17	0	0	0	0	0	0	5375	17	17	0	5391
2003–04	0	0	0 ^f	0	0 ^h	4874	0 ^g	4874	0	226 ^e	226	0	20	20	0	0	0	4874	247	247	0	5120
2004–05	0	0	0	0	0	5214	0	5214	0	35	35	0	0	0	0	0	0	5214	35	35	0	5248
2005–06	0	0	0	0	0	5302	0	5302	0	6	6	0	0	0	0	0	0	5302	6	6	0	5308
2006–07	0	0	0	0	0	5230	0	5230	0	4	4	0	0	0	0	0	0	5230	4	4	0	5234
2007–08	0	0	0	0	0	5211	0	5211	0	23	23	0	0	0	0	0	0	5211	23	23	0	5234
2008–09	0	0	0	0	2	5015	0	5017	11	213	225	0	<1	<1	0	0	0	5029	213	213	0	5242
2009–10 ⁱ	0	0	0	0	0	3931	0	3931	0	161	161	0	0	0	0	0	0	3931	161	161	0	4091
2010–11 ^j	0	0	0	0	0	3872	0	3872	0	85	85	1	0	1	0	0	0	3872	85	85	0	3958
2011–12	0	0	0	0	0	4485	0	4485	0	58	58	0	0	0	0	0	0	4485	58	58	0	4543
2012–13	0	0	<1	0	0	4198	0	4198	0	341	341	0	0	0	0	0	0	4198	341	341	0	4539
2013–14	0	0	0	0	0	5039 ^k	11	5050	0	369	369	0	0	0	0	0	0	5039	380	380	0	5420
2014–15	0	0	0	0	0	4948 ^l	0	4948	0	572 ^m	572	0	0	0	0	0	0	4948	571	571	0	5519
2015–16	0	0	0	0	0	4899 ⁿ	0	4899	0	554	554	0	180 ^o	180	0	0	0	4899	734	734	0	5633
2016–17	0	0	0	0	0	4683	0	4683	0	566 ^p	566	0	84 ^q	84	0	0	0	4683	650	650	0	5334
2017–18	0	0	0	0	0	5130 ^r	0	5130	0	1029 ^s	1029	0	0	0	0	0	0	5130	1029	1029	0	6159
2018–19	0	0	0	0	0	5294 ^t	0	5294	0	761	761	0	20 ^u	20	0	0	0	5308	766	766	0	6074
2019–20	0	0	0	0	0	4570 ^v	0	4570	0	833	833	0	26 ^w	26	0	0	0	4586	844	844	0	5429

Note that 'RTMP' refers to the Real Time Monitoring Program.

^a Note that a further 700 t of Australian quota was 'frozen' (not allocated) in 1990–91.

^b 1997–98 and 1998–99 WA and SA non-farm catches are included in SA pole and purse seine catch, and in 1999–00 and 2000–01 WA longline catch is included in SA longline due to confidentiality guidelines.

^c 1997–98 to 1998–99 NSW pole and purse seine catches are included in NSW longline catch due to confidentiality guidelines.

^d 1997–98 and 1998–99 TAS troll catches are included in TAS longline, and in 1999–00, 2000–01 and 2001–02 TAS longline catch is included in NSW longline due to confidentiality guidelines.

^e 2001–02 and 2003–04 NSW longline catch also includes QLD longline catch due to confidentiality guidelines.

^f 2002–03 and 2003–04 WA longline catch is included in NSW longline due to confidentiality guidelines.

^g 2003–04 SA longline catch is included in NSW longline due to confidentiality guidelines.

^h 2003–04 additional SA purse seine catch that did not go into farm cages is included in SA farm cages catch due to confidentiality guidelines.

ⁱ Year 1 of the 2009–11 fishing season (2009–10).

^j Year 2 of the 2009–11 fishing season (2010–11).

^k Includes <1t of trolling.

^l Includes <1t of pole-and-line.

^m Includes <5t of rod-and-reel around NSW and rod-and-reel and longline around Tasmania, due to confidentiality constraints.

ⁿ Includes <5t of pole-and-line and rod-and-reel, due to confidentiality constraints.

^o Includes <1t of trolling, due to confidentiality constraints.

^p Includes <1t of trolling, due to confidentiality constraints.

^q Includes <1t of trolling, rod-and-reeling and gillnetting, due to confidentiality constraints.

^r Includes pole-and-line, due to confidentiality constraints.

^s Includes longlining and rod-and-reeling off Tasmania, due to confidentiality constraints.

^t Includes pole-and-line, due to confidentiality constraints.

^u Includes trolling and rod-and-reeling, due to confidentiality constraints.

^v Includes pole-and-line, due to confidentiality constraints

^w Includes trolling and rod-and-reeling, due to confidentiality constraints

2 Catch and effort

2.1 Purse seine fishery

In 2019–20, 84.1% of the Australian catch of SBT was taken by purse seine off SA for farm operations. The remainder was taken by longline off NSW; pole-and-line off South Australia; and, longline, trolling and rod-and-reel off Tasmania. Australian catch by gear and state from the 1988–89 to the 2019–20 fishing season is shown in Table 1. Catch by fishing season with number of vessels and vessel search hours are shown in Appendix 3. The Australian catch of SBT for the calendar years 2019 and 2020 is mapped in Figure 1 and Figure 2, respectively.

2.1.1 Nominal CPUE

Nominal catch-per-unit-effort (CPUE) indices for the Australian surface and longline fisheries are not viewed as indicative of stock status. The farm operations use purse seines to catch SBT, with assistance from bait vessels and spotter planes. Australia does not consider these data suitable for interpreting catch rates as an index of abundance.

2.1.2 Size composition

In the SA purse seine fishery there has been reduced competition for SBT among fishers following the introduction of individual transferable quotas (ITQs). Starting in the late 1980s, the fishery targeted small (~80 cm) SBT, which were previously taken in bulk for canning (Caton et al. 1995). However, the size range of SBT taken has been variable since then. Closure of Australian canneries resulted in an increase in the average length of SBT landed for fresh-chilled export. As the farming component of the fishery increased in the 1990s, the average length of SBT landed in SA decreased from a peak of 103.2 cm in 1995 to a low of 81.9 cm in 2011 (Table 2). This is primarily due to selective targeting of schools to catch the best sized fish for farming. The average length of SBT landed in SA in 2019–20 was 87.8 cm.

2.2 Longline fishery

The catch of SBT by the Australian longline fleet off the east coast has grown in recent years from 58 t in 2011–12 to 844 t in 2019–20. Approximately 95% of the Australian longline caught SBT are landed or processed in five main locations: Ulladulla, Bermagui, Eden, Sydney and Mooloolaba. Current in port sampling programs are mainly targeted at tropical tunas, however SBT length frequency data are also collected from the major ports during the months May – October (n=1767 in 2019–20).

As the catch has increased questions have been raised about whether it is necessary to collect otoliths for estimation of the age composition of the catch of this fishery (for use in the SBT operating models), and if so, how many otoliths are required and how will length-at-age data be used to estimate catch-at-age with required level of precision. The size range of SBT caught in the longline fishery (~110–180 cm fork length) is larger than those caught by the surface fishery (~80–110 cm) so length-at-age data currently collected by Australia cannot be applied to the longline fishery.

Australia engaged CSIRO in 2017 to evaluate the direct ageing requirements for the Australian longline fishery. The research concluded that based on the size range of SBT caught by the longline fishery a minimum sample size of 300 otoliths (~20 fish per 5-cm length class) for direct ageing per fishing season would be sufficient. It was noted that flexibility in the sampling program is important if the size range of SBT in the catch changes over time. The aim of the sampling program is to provide a representative estimate of length at age by collecting otoliths from the full range of sizes caught while maintaining sufficient numbers of otoliths in each size class to achieve the target level of precision. Australia is currently investigating options for the ongoing collection of otoliths and other data, however travel restrictions due to the COVID-19 pandemic have limited collection opportunities in recent seasons.

2.2.1 Nominal CPUE

Australian longliners generally target more than one species in the fishing season and the targeted effort (number of hooks targeting SBT) is not distinguishable from logbooks. For information, nominal SBT CPUE for all Australian longline effort is provided at Appendix 4.

2.2.2 Size composition

Since the late 1980s, the average length of SBT landed in NSW has varied considerably because of the varying contribution of longline and sporadic surface catches to the overall catch levels. However, longline-caught SBT off NSW have, in general, been considerably larger than SBT previously taken in this fishery by purse seine in South Australia (Figure 4). Similarly, the size trends in the Tasmania area (TAS) of the fishery reflect the change from trolling to longlining operations, which target larger fish, in 1993.

The percentage representation by length in the winter catches of Japanese longliners off eastern TAS from 1988 to 1997 show modal changes through time (Figure 5). Modes that represent juvenile age classes were less visually evident in 1988 and 1989. Sample sizes were < 1000 in these years which may contribute to the lack of modal clarity. Modes at approximately 90 cm, 104 cm and 118 cm were visually evident in the period between 1991 and 1997. Progression of these modes into size classes >130 cm was also visually evident over this period. The absence of clear modes in the size range of 130 cm to 150 cm in the period 1988 to 1993 is consistent with intensive removals of small SBT in the early 1980s by Australia's surface fishery.

A reduced representation of SBT <105 cm was visually evident in 1994, then of SBT <120 cm in 1995, and SBT <135 cm in 1996. This coincided with a reduced abundance of small SBT (<100 cm) in the TAS troll fishery in 1994 and 1995, and the failure of the troll fishery in 1996 and 1997. The Japanese and joint-venture longline operations in the AFZ ceased operations in 1997.

Figure 1 Australian SBT catch in the 2019 calendar year. Note that catch is centred in each grid square.




Figure 2 Australian SBT catch in the 2020 calendar year. Note that catch is centred in each grid square.




Table 2 Average fork length (cm) of SBT landed in each Australian state, 1989 to 2020

Calendar Year	Western Australia ^a	South Australia ^a	Tasmania	NSW	Joint-venture
1989	65.4	88.8	–	–	–
1990	65.6	89.3	96.0	112.8	–
1991	67.2	95.5	94.9	154.8	114.5
1992	66.1	97.0	93.4	109.2	108.0
1993	65.2	101.1	99.7	117.8	116.5
1994	–	97.4	125.5	121.3	124.8
1995	–	103.2	127.9	125.0	125.0
1996	–	102.7	132.7	139.7	–
1997	–	97.7	133.2	134.6	–
1998	–	94.9	134.5	136.1	–
1999	–	97.6	134.2	138.5	–
2000	–	97.0	–	154.3	–
2001	154.3	98.1	–	149.7	–
2002	–	98.4	–	159.9	–
2003	–	98.7	–	154.1	–
2004	–	93.6	–	161.9	–
2005	–	91.0	–	161.7	–
2006	–	90.7	–	154.1	–
2007	–	94.0	–	150.5	–
2008	–	93.9	–	166.7	–
2009	–	97.3	159.9	149.6	–
2010	–	94.4	–	146.7	–
2011	–	89.1	147.5	149.1	–
2012	–	93.0	–	167.0	–
2013	–	96.1	–	146.6	–
2014	–	94.7	–	155.2	–
2015	–	92.5	–	140.3	–
2016	–	96.4	–	142.9	–
2017	–	93.4	134.1	140.4	–
2018	–	90.6	133.5	139.5	–
2019	–	87.8	129.6	140.7	–
2020	–	–	132.5	141.4	–

^a Lengths are reported by calendar year, except for Western Australia, which are by financial year (e.g. 1998 represents the financial year 1998–99) and South Australia, which are by fishing season, to cover the summer season

Australia's 2019–20 Southern Bluefin Tuna Fishing Season




Figure 3 Length frequency of SBT purse seine catch in Australian waters raised to total catch, 2012–13 to 2019–20 fishing seasons (source: tow cage size monitoring database)

Australia's 2019–20 Southern Bluefin Tuna Fishing Season




Figure 4 Length frequency histograms for retained SBT longline catch in Australian waters raised to total catch, 2012–13 to 2019–20 quota years (source: AFMA observer data and processor monitoring data)




Figure 5 Length frequency of SBT measured by Australian observers on Japanese longliners fishing in the Tasmania region, 1988 to 1997. The data include small SBT tagged and released by observers.

3 Fleet size and distribution

In 2019–20, a total of 39 commercial fishing vessels landed SBT (or transferred to farm cages) in Australian waters.

3.1 South Australia (SA)

The one- to five-year-old SBT, which school from late spring to autumn in surface waters of the eastern Great Australian Bight, SA, were fished by seven purse seiners during 2019–20, with various live bait, pontoon-towing and feeding vessels also involved. The majority of the purse seine fishing commenced in mid-December 2019 and finished in mid-March 2020. Two vessels conducted pole-and-line operations in the waters off SA in the 2019–20 season.

3.2 Western Australia (WA)

No vessels caught SBT off WA in 2019–20.

3.3 New South Wales (NSW)

In 2019–20, 22 longliners reported landing both older juvenile and adult SBT in deeper waters off NSW. Two of these vessels also fished waters around Tasmania and two of these vessels also fished waters off Queensland.

3.4 Tasmania (TAS)

In 2019–20, two longliners caught SBT off Tasmania. They also operated in waters off NSW. Three vessels conducted rod-and-reel operations catching SBT off Tasmania. Four vessels conducted trolling operations catching SBT off Tasmania.

3.5 Queensland (QLD)

Three vessels caught SBT off Queensland in 2019–20.

4 Research and monitoring to improve estimates of attributable catch

4.1 Recreational fishing

Recreational fishing for SBT occurs primarily off south-east TAS, NSW, SA, and western VIC. There is also some catch of small SBT off south-west WA. Recreational fishing for SBT is managed by the relevant states. States that have a recreational fishing bag limit (number of fish that can be retained) for SBT include SA, VIC, TAS and NSW. SA also has a limit on the number of SBT taken per boat. Retention of SBT by recreational fishers is banned in QLD.

Tracey et al. (2016) details a study examining post-release survival and physiological stress in SBT captured recreationally. This study estimated a post-release survival rate of 86.6% and indicates that SBT captured recreationally are likely to survive when released. However, this result is a best-case scenario for recreational fishing using proper handling methods and it is unclear how these results relate to commercial longline post-survival rates.

At CCSBT24, Australia advised that it had set aside an amount of 250 t to account for recreational catch. Australia noted that it would also settle agreement with relevant states on resource sharing and management, implement a program to educate recreational fishers on SBT and fish handling practices.

4.1.1 National estimate of recreational SBT catch

In 2013, Australia commenced a project with the relevant State jurisdictions to develop a methodology to obtain robust estimates of the total Australian recreational catch of SBT (Moore et al. 2015). The project incorporated elements of previous surveys, including Tasmanian off-site and Victorian on-site surveys. The project reviewed potential methodologies, conducted initial survey design work and tested on-site surveys in SA locations, from January to August 2014.

A national survey of recreational fishing for SBT that commenced in 2018 has now been finalised (Tracey et al. 2020) and the survey report can be found at https://www.imas.utas.edu.au/_data/assets/pdf_file/0003/1331796/National-Survey-of-Rec-Fishing-for-SBT-in-Australia.pdf. The survey estimated recreational catch was 270 t for the survey year. Australia advised the CCSBT in 2020 that it would set aside 5% of its CCSBT allocation annually against recreational catch, which at the time was around 308 t. The additional amount set aside allows for fluctuation between years in the recreational fishery, plus some room for growth in recreational catch.

4.1.2 Variation in recreational catch

Anecdotal evidence suggests that the recreational catch of SBT is can vary between years and regions. The mechanisms driving this variability are not well understood but likely include factors such variation in the currents and oceanographic conditions. These environmental factors also likely affect the size of the fish that are available, as variation in the size of the SBT taken by recreational fishers has been documented both within and between years (e.g. Forbes et al. 2009; Tracey et al. 2013).

Fisher behaviour also varies between regions and seasons and is likely to influence the level of fishing for SBT. In years where the fish are further offshore there is likely to be less effort to fish. In contrast, when the fish occur close to shore they may be available to a wider group of fishers. Fishers may also be motivated to fish (or not to fish) based on other factors such as the size of the fish being taken and the perceived abundance of fish in that year.

4.1.3 Tag releases

The number of reported SBT tag releases by Australian recreational fishers is provided in Table 3. These data do not include recreational tag releases using CCSBT tags. There has been an increase in reported tag and release activity for SBT since 2005, particularly in SA. The trends in SBT tag-releases are difficult to interpret; tagging is influenced by various factors, including the abundance and distribution of SBT, angler participation rates (fishing effort), angler attitudes and behaviour, the condition of fish, the availability of tags and management measures (e.g. bag limits) in relation to fish size and catch rates.

Table 3 Reported tag releases of SBT by Australian recreational fishers, 1990 to 2020

Year	Percentage of total releases					No. of Releases
	TAS	NSW	SA	VIC	WA	
1990	7.1	0.0	14.3	57.1	21.4	14
1991	5.8	44.7	23.2	25.6	0.7	293
1992	5.6	18.5	48.1	24.1	3.7	54
1993	3.5	6.9	87.9	0.0	1.7	231
1994	0.0	14.3	76.2	0.0	9.5	63
1995	0.0	25.0	25.0	0.0	50.0	12
1996	0.0	25.8	74.2	0.0	0.0	159
1997	0.0	3.7	87.0	0.0	9.3	54
1998	4.8	0.8	52.5	40.6	1.3	377
1999	0.0	0.0	70.9	29.1	0.0	117
2000	0.0	0.4	99.1	0.4	0.0	224
2001	38.8	1.9	23.8	30.6	5.0	160
2002	39.4	2.6	25.9	25.4	6.7	193
2003	14.3	0.0	71.4	0.0	14.3	21
2004	31.8	0.0	45.5	0.0	22.7	22
2005	0.0	2.1	93.7	0.0	4.2	96
2006	7.2	0.2	84.7	0	7.9	584
2007	29.3	0.1	66.2	1.3	3.1	1303
2008	1.0	0	96.6	0.5	1.9	1006
2009	9.0	3.3	83.9	1.7	2.2	2044
2010	5.2	0.5	85.0	4.6	4.6	3501
2011	4.3	15.9	75.7	3.5	0.6	3351
2012	36.5	5.4	40.6	14.3	3.3	1042
2013	12.8	0.3	80.0	6.6	0.3	986
2014	22.9	3.0	59.5	14.1	0.5	2288
2015	16.4	1.0	72.3	9.7	0.6	2106
2016	35.8	0.6	55.3	8.1	0.2	2842
2017	29.3	2.2	58.9	8.9	0.8	1179
2018	18.5	2.8	56.9	21.0	0.7	1402
2019	28.9	0.5	46.2	23.2	1.2	1095
2020	42.1	0.3	32.8	24.2	0.7	1174

Source: NSW DPI Game Fish Tagging Program

4.2 Discards in the commercial fishery

4.2.1 Describe the various sources of information and data used in calculating the estimates

Purse-seine sector

Discarding of fish other than live release (immediately after encirclement or as a live release from tow cage) is not permitted in the purse seine sector. All dead fish encountered during purse seine operations, towing or transfers must be recorded in logbooks and accounted for under the quota system. To validate logbooks, observers and at sea inspections are used. In the 2019–20 fishing season, observers monitored 9.9% of purse seine sets where fish were retained for the farm sector and 10.3% of the estimated SBT catch. 100% of all transfers were monitored by Australian Fisheries Management Authority (AFMA) representatives in Port Lincoln. During 2019–20, no discarding of SBT, excluding observed live releases noted in the observer section of the report, was reported in logbooks collected in the purse-seine fishery in the Great Australian Bight.

Longline sector

During the period from May to October, the waters off the east coast of New South Wales (Australia) become an area of significant interaction between the Southern Bluefin Tuna Fishery (SBTF) and the Eastern Tuna and Billfish Fishery (ETBF). While the ETBF is a multi-species fishery, the SBT Fishery is a single species fishery that requires operators to hold SBT quota that is nominated to their boat in order to take the species.

To address the risk of interactions with SBT for vessels who do not hold quota, AFMA institutes restricted access areas in the ETBF annually. These arrangements require ETBF operators to have minimum SBT quota holdings and a fully operational electronic monitoring system installed in order to operate in designated areas of the ETBF where SBT are likely to interact with longline fishing gear.

The location and timing of the SBT zone is determined by analysing information from a range of sources; sea surface temperatures, landing data, VMS data and advice from industry. This information provides AFMA with an indication of the area where SBT are likely to occur.

The Southern Bluefin Tuna Fishery Management Plan 1995 outlines the rules relating to the fishery. Statutory fishing rights issued under the Plan state the following in relation to discarding of SBT in the longline sector.

The conditions for retaining and discarding SBT in the longline sector are outlined below (see [https://www.afma.gov.au/sites/default/files/southern_bluefin_tuna_sfr_general_conditions_2019 - 20_season.pdf](https://www.afma.gov.au/sites/default/files/southern_bluefin_tuna_sfr_general_conditions_2019_-_20_season.pdf) for full conditions)

When fishing in the SBT Zone using the longline method the holder must:

- a) on first entry (which includes being present or fishing in) to the SBT Zone hold at least 1 uncaught Southern Bluefin Tuna Statutory Fishing Right nominated to the boat nominated to this concession;*
- b) on every subsequent entry to the SBT Zone continue to hold at least 1 uncaught Southern Bluefin Tuna Statutory Fishing Right nominated to the boat nominated to this concession;*
- c) ensure all dead SBT are landed (retained) except those that are damaged by sharks to the point that they are not fit for human consumption;*
- d) ensure live SBT are either landed (retained) or released (providing they are in a live and vigorous state).*
- e) ensure any SBT that are to be released or discarded must be done so at the fish door in full view of the e-monitoring system. SBT damaged by sharks must have the damaged portion facing the cameras. SBT that are undamaged and dead when discarded or live fish that are gaffed and released will be deducted from the concession holder's quota. The amount of quota decremented will be determined by the average weight of SBT retained for that trip.*
- f) ensure all SBT, whether retained, released or discarded are reported in the logbook.*
- g) carry an AFMA observer when notified by AFMA.*

E-Monitoring

The baseline (e-monitoring) audit rate for all fisheries is a minimum 10% of shots per boat and a minimum of one shot per drive for each boat per month. The analyses include analysis of full catch composition for each shot selected for review. Catch composition, discards (including life status) and interactions with protected species on audited shots will be compared to logbook records with discrepancies flagged and reported. The focus of e-monitoring is on fishing activities. However, if behaviour that contravenes Australian or International law is observed in the process of viewing footage, it will be referred to the AFMA Compliance team for investigation.

4.2.2 Describe the method applied for estimating the catch

In 2020, e-monitoring observed 11.0% of longline hook effort in the ETBF during the months and in the areas of the SBT migration through that fishery. For the fishery as a whole, 9.8% of hooks were observed in 2020. The observed (via e-monitoring) total catch number of SBT was 1,086 individuals, of which 1,044 were retained, 42 fish were discarded (19 of which were reported as released alive, 9 were dead or damaged and 14 of which were reported as having an undetermined life status); none were tagged. Raised mortality from these figures is 91.8 fish. Average whole weight of SBT retained in 2020 was 50.4kg.

4.2.3 Provide the resulting estimated catch

Estimated shark damaged discards of SBT from the commercial fleet in 2020 is 4.6 tonnes, assuming the average weight of shark damaged fish is the same as landed fish (50.4kg).

4.3 Customary and/or traditional fishing

Not applicable

4.4 Artisanal fishing

Not applicable

5 Other relevant information

5.1 Stereo-video commercial trial

In 2011, the commercial trial of stereo-video technology in the SBTF was used to record SBT from eight transfers made from three cages. In total, 23,018 SBT were counted from the stereo-video recordings and overall stereo-video technology was demonstrated to be robust under commercial operating conditions. Technical details of the commercial trial were reported to the CCSBT Compliance Committee Meeting in 2011 (Anon. 2011; CCSBT-CC/1110/11). Following the first trial, the Australian Government adopted a policy that stereo video would not be implemented in Australian tuna farms until the technology was 'cost effective and fully automated'.

General work examining the automation of measurements of fish using stereo-video was reported in Shaifait et al. (2017)

(<https://academic.oup.com/icesjms/article/74/6/1690/3056949>). Copies of this paper were provided to CCSBT 24.

In 2020, Australia advised the CCSBT that it would again trial the cost effectiveness and accuracy of fully automated stereo systems in situ in Australia's tuna farms. Australia will report again to the CCSBT Annual Meeting this year on progress towards this trial.

Appendix A: SBT fishing season dates 1988-89 to 2019-20

Quota Year	Start Date	End Date
1988–89	1 Oct 1988	30 Sep 1989
1989–90	1 Oct 1989	30 Sep 1990
1990–91	1 Oct 1990	30 Sep 1991
1991–92	1 Oct 1991	31 Oct 1992
1992–93	1 Nov 1992	31 Oct 1993
1993–94	1 Nov 1993	31 Oct 1994
1994–95	1 Nov 1994	15 Dec 1995
1995–96	16 Dec 1995	15 Dec 1996
1996–97	16 Dec 1996	30 Nov 1997
1997–98	1 Dec 1997	30 Nov 1998
1998–99	1 Dec 1998	30 Nov 1999
1999–00	1 Dec 1999	30 Nov 2000
2000–01	1 Dec 2000	30 Nov 2001
2001–02	1 Dec 2001	30 Nov 2002
2002–03	1 Dec 2002	30 Nov 2003
2003–04	1 Dec 2003	30 Nov 2004
2004–05	1 Dec 2004	30 Nov 2005
2005–06	1 Dec 2005	30 Nov 2006
2006–07	1 Dec 2006	30 Nov 2007
2007–08	1 Dec 2007	30 Nov 2008
2008–09	1 Dec 2008	30 Nov 2009
2009–11 ^a	1 Dec 2009	30 Nov 2011
2011–12	1 Dec 2011	30 Nov 2012
2012–13	1 Dec 2012	30 Nov 2013
2013–14	1 Dec 2013	30 Nov 2014
2014–15	1 Dec 2014	30 Nov 2015
2015–16	1 Dec 2015	30 Nov 2016
2016–17	1 Dec 2016	30 Nov 2017
2017–18	1 Dec 2017	30 Nov 2018
2018–19	1 Dec 2018	30 Nov 2019
2019–20	1 Dec 2019	30 Nov 2020

^a Note that 2009–11 was a two-year season

Appendix B: Purse seine fishing season duration

Quota Year	First Day of Season	Last Day of Season	1st Day	50%	75%	90%	Last Day	Duration
1993–94	1 Nov 93	31 Oct 94	49	113	127	138	175	127
1994–95	1 Nov 94	15 Dec 95	36	106	133	160	410 ^a	375 ^a
1995–96	16 Dec 95	15 Dec 96	1	67	87	131	365	365
1996–97	16 Dec 96	30 Nov 97	2	66	85	95	141	140
1997–98	1 Dec 97	30 Nov 98	19	67	84	98	364	346
1998–99	1 Dec 98	30 Nov 99	10	52	73	78	113	104
1999–00	1 Dec 99	30 Nov 00	4	56	65	79	118	115
2000–01	1 Dec 00	30 Nov 01	4	60	80	88	97	94
2001–02	1 Dec 01	30 Nov 02	9	61	75	80	121	113
2002–03	1 Dec 02	30 Nov 03	11	60	82	97	116	106
2003–04	1 Dec 03	30 Nov 04	9	66	87	102	115	107
2004–05	1 Dec 04	30 Nov 05	5	61	83	98	119	115
2005–06	1 Dec 05	30 Nov 06	18	70	92	99	358	341
2006–07	1 Dec 06	30 Nov 07	1	74	93	104	125	125
2007–08	1 Dec 07	30 Nov 08	10	58	91	94	99	90
2008–09	1 Dec 08	30 Nov 09	3	76	103	113	130	128
2009–11 ^b	1 Dec 09	30 Nov 10	3	52	69	78	84	82
2009–11 ^c	1 Dec 10	30 Nov 11	22	61	87	100	356	335
2011–12	1 Dec 11	30 Nov 12	22	71	85	99	110	89
2012–13	1 Dec 12	30 Nov 13	21	57	71	79	102	82
2013–14	1 Dec 13	30 Nov 14	18	58	63	64	69	52
2014–15	1 Dec 14	30 Nov 15	38	72	89	94	112	75
2015–16	1 Dec 15	30 Nov 16	19	62	72	75	86	68
2016–17	1 Dec 16	30 Nov 17	21	47	57	66	90	70
2017–18	1 Dec 17	30 Nov 18	15	67	80	92	113	99
2018–19	1 Dec 18	30 Nov 19	22	69	84	90	108	87
2019–20	1 Dec 19	30 Nov 20	15	58	85	102	107	93

'1st Day' = Day of First Capture, '50%', etc denote the day of the season on which that percentage of the catch had been taken, 'Last Day' = the Day of Last Capture.

^aFigures greater than 365 days because the season dates changed and extended this season for longer than one year.

^bYear 1 of the 2009–11 season (2009–10)

^cYear 2 of the 2009–11 season (2010–11)

Appendix C: Australian surface catch for farm operations, 1994-95 to 2019-20


Season	Estimated catch (t)	Actual catch (t)	Catcher vessels	Vessel search hours	Sets	No. 1° squares fished
1994-95	2179	2009	5	526	104	5
1995-96	2859	3442	6	631	89	11
1996-97	3134	2505	7	769	118	13
1997-98	3916	3629	7	671	143	8
1998-99	4418	4991	7	972	129	3
1999-00	4746	5131	8	764	107	5
2000-01	5100	5162	8	799	129	2
2001-02	5400	5234	7	1309	159	3
2002-03	5188	5375	7	1276	150	5
2003-04	5299	4874	6	1202	160	4
2004-05	5225	5215	8	1168	139	4
2005-06	5463	5302	7	1304	156	6
2006-07	5091	5230	6	1459	160	8
2007-08	4530	5211	7	1217	134	2
2008-09	4348	5017	7	1156	139	7
2009-11 ^a	3323	3931	6	417	78	3
2009-11 ^b	3840	3872	5	835	106	5
2011-12	4328	4485	5	1150	156	7
2012-13	4039	4198	5	1021	110	11
2013-14	4381	5029	6	752	101	4
2014-15	4789	4950	6	1235	154	6
2015-16	4826	4896	6	1076	124	6
2016-17	4036	4683	6	1004	109	5
2017-18	4920	5123	7	1137	191	6
2018-19	4750	5291	7	1366	154	9
2019-20	4224	4568	7	1248	142	12

Note that estimated catch is derived from logbook data while actual catch is derived from landing data.


^aYear 1 of the 2009–11 season (2009–10)

^bYear 2 of the 2009–11 season (2010–11)

Appendix D: Nominal CPUE for all Australian longliners, 2019 and 2020 calendar years



Australia's 2019–20 Southern Bluefin Tuna Fishing Season



Appendix E: Summary of observed catch and effort by year and sector

Country	Year	Sector	Observers Deployed	Sea Days	Sets/Tows Observed	Observed Vessels	Observed Effort (% , units)	Observed Catch (% , units)	Total Cost
Australia	2002–03	Purse Seine ^a	N/A	47	24		11% (sets)	11% (est. total weight)	60,000 (A\$)
Australia	2002–03	Towing ^a	N/A	19	1		2.6% (tows)		(included above)
Australia	2002	East Coast Longline	17	323	198		14.4% (hooks)	35.5% (no. retained catch)	NA
Australia	2002	West Coast Longline	N/A	N/A	N/A		N/A (hooks)	N/A (no. retained catch)	NA
Australia	2003–04	Purse Seine ^a	2	27	21		13% (sets)	12.8% (est. total weight)	60,000 (A\$)
Australia	2003–04	Towing ^a	2	30	2		5.6% (tows)		(included above)
Australia	2003	East Coast Longline	10	242	168		14.9% (hooks)	55.2% (no. retained catch)	303,000 (60,000 A\$ SBT component)
Australia	2003	West Coast Longline	4	72	54		2.0% (hooks)	4.5% (no. retained catch)	42,247 (A\$)
Australia	2004–05	Purse Seine ^a	2	36	15		11.2% (sets)	8.5% (est. total weight)	60,000 (A\$)
Australia	2004–05	Towing ^a	2	24	2		5.7% (tows)		(included above)
Australia	2004	East Coast Longline	11		68		11.7% (hooks)	5.4% (no. retained catch)	966,000 (150,000 A\$ SBT component)
Australia	2004	West Coast Longline			59		3.9% (hooks)	0% (no. retained catch)	57,384(A\$)
Australia	2005–06	Purse Seine ^a	2	47	14		9.2% (sets)	10.1% (est. total weight)	78,000 (A\$)
Australia	2005	East Coast Longline	14		128		37.5% (hooks)	62.8% (no. retained catch)	723,289 (160,000 A\$ SBT component)
Australia	2005	West Coast Longline			47		9.1% (hooks)	(no observed catch)	0

Australia's 2019–20 Southern Bluefin Tuna Fishing Season

Country	Year	Sector	Observers Deployed	Sea Days	Sets/Tows Observed	Observed Vessels	Observed Effort (% , units)	Observed Catch (% , units)	Total Cost
Australia	2006–07	Purse Seine ^a	2	50	9		5.6% (sets)	12.1% (est. total weight)	NA
Australia	2006–07	Towing ^a	2	41	2		6.5% (tows)		NA
Australia	2006	East Coast Longline	20		138		22.1% (hooks)	88.9% (no. retained catch)	NA
Australia	2006	West Coast Longline	1		8		17.4% (hooks)	(no observed catch)	NA
Australia	2007–08	Purse Seine ^a	2	19	16		11.8% (sets)	5.6% (est. total weight)	68,000 (A\$)
Australia	2007–08	Towing ^a	2	38	2		6.0% (tows)		(included above)
Australia	2007	East Coast Longline	17		156		30.2% (hooks – SBT Area)	23.2% (no. retained catch)	180,000 (A\$)
Australia	2007	West Coast Longline			10		1.9% (hooks)	No SBT caught	15,589 (A\$)
Australia	2008–09	Purse Seine	2	27	11 (fish retained) 8 (aborted)	3	7.9% (sets, fish retained)	15.3% (est. total weight)	77,215 (A\$)
Australia	2008–09	Towing	1	15	1	1	3.2% (tows)		(included above)
Australia	2008	East Coast Longline	31		676		47.9% (hooks – SBT Area)	34% (no. retained catch)	694,500 (A\$ – 08/09 fin year)
Australia	2008	West Coast Longline	3		25		16.7% (sets)	No SBT caught	16,800 (A\$ – 08/09 fin year)
Australia	2009–11 ^b	Purse Seine	1	3	7 (fish retained) 1 (aborted)	2	9.0% (sets, fish retained)	13.5% (est. total weight)	95,392 (A\$)
Australia	2009–11 ^b	Towing	1	27	1	1	4.2% (tows)		(included above)
Australia	2009	East Coast Longline	20		400		17.2% (hooks – SBT Area)	23% (no. retained catch)	332,562 (A\$ – 09/10 fin year)
Australia	2009	West Coast Longline	2		31		8.2% (sets)	No SBT caught	21,019 (A\$ – 09/10 fin year)
Australia	2009–11 ^c	Purse Seine	2	49	21 (fish retained) 11 (aborted)	2	19.8% (fish retained)	12.2% (est. total weight)	48,830 (A\$)
Australia	2009–11 ^c	Towing	2	22	1	1	3.7% (tows)		(included above)
Australia	2010	East Coast Longline	16		65		7.7% (hooks – SBT Area)	20.1% (no. retained catch)	417,240 (A\$ – 10/11 fin year)
Australia	2010	West Coast Longline	1		10		2.5% (hook effort)	No SBT caught	14,533 (A\$ – 10/11 fin year)

Australia's 2019–20 Southern Bluefin Tuna Fishing Season

Country	Year	Sector	Observers Deployed	Sea Days	Sets/Tows Observed	Observed Vessels	Observed Effort (% , units)	Observed Catch (% , units)	Total Cost
Australia	2011–12	Purse Seine	1	17	17 (fish retained) 2 (aborted)	1	11.1% (fish retained)	13.8% (est. total weight)	45,000 (A\$)
Australia	2011–12	Towing	1	13	1	1	3.4% (tows)		(included above)
Australia	2011	East Coast Longline	9		76		9.6% (hooks – SBT Area)	17.7% (no. retained catch)	\$750,000 (A\$)
Australia	2011	West Coast Longline	1		4		1.7% (hook effort)	No SBT caught	\$11,500 (A\$)
Australia	2012–13	Purse Seine	2	30	14 (fish retained) 1 (aborted)	2	12.7% (fish retained)	13.9% (est. total weight)	\$75,000 (A\$)
Australia	2012–13	Towing	2	26	2	2	3.8% (tows)		(included above)
Australia	2012	East Coast Longline	9		50		6.2% (hooks – SBT Area)	16.1% (no. retained catch)	\$800,000 (A\$)
Australia	2012	West Coast Longline	3		61		17.9% (hook effort)	No SBT caught	\$90,000 (A\$)
Australia	2013–14	Purse Seine	2	17	16 (fish retained) 1 (aborted)	2	17.0% (fish retained)	21.9% (est. total weight)	na
Australia	2013–14	Towing	1	9	1	1	4% (tows)		
Australia	2013	East Coast Longline	10		87		10.4% (hooks – SBT Area)	19.5% (no. retained catch)	na
Australia	2013	West Coast Longline	0		0		0%	No observer coverage	
Australia	2014–15	Purse Seine	1	17	14 (fish retained)	1	9.1% (fish retained)	19.9% (est. total weight)	na
Australia	2014–15	Towing	1	20	1	1	4% (tows)		
Australia	2014	East Coast Longline	5		24		3.1% (hooks – SBT Area)	4.5% (no. retained catch)	na
Australia	2014	West Coast Longline	2		27		9.1%	31.8% (no. retained catch)	
Australia	2015–16	Purse Seine	2	15	25	2	18.9% (fish retained)	20.2% (est. total weight)	na
Australia	2015–16	Towing	2	21	2	2	7.1% (tows)		
Australia	2015	East Coast Longline			330		5.9% (hooks – SBT Area)	6.5% (no. retained catch)	na
Australia	2015	West Coast Longline			19		7.2%	No observed catch	na

Australia's 2019–20 Southern Bluefin Tuna Fishing Season

Country	Year	Sector	Observers Deployed	Sea Days	Sets/Tows Observed	Observed Vessels	Observed Effort (% , units)	Observed Catch (% , units)	Total Cost
Australia	2016–17	Purse Seine	2	11	20	2	18.3% (fish retained)	16.8% (est. total weight)	na
Australia	2016–17	Towing	2	18	2	2	9.1% (tows)		
Australia	2016	East Coast Longline			397		9.3% (hooks – SBT Area)	12.1% (no. retained catch)	na
Australia	2016	West Coast Longline			24		10.2%	No observed catch	na
Australia	2017–18	Purse Seine	2	37	40	2	20.9% (fish retained)	19.0% (est. total weight)	na
Australia	2017–18	Towing	1	20	1	2	3.4% (tows)		
Australia	2017	East Coast Longline			527		9.0% (hooks – SBT Area)	9.7% (no. retained catch)	na
Australia	2017	West Coast Longline			32		11.7%	No observed catch	na
Australia	2018–19	Purse Seine	2	35	22	2	14.3% (fish retained)	14.5% (est. total weight)	na
Australia	2018–19	Towing	2	25	2	2	7.7% (tows)		
Australia	2018	East Coast Longline			488		11.5% (hooks – SBT Area)	13.8% (no. retained catch)	na
Australia	2018	West Coast Longline			36		13.0%	No observed catch	na
Australia	2019–20	Purse Seine	2	34	14	2	9.9% (fish retained)	10.3% (est. total weight)	na
Australia	2019–20	Towing	2	21	2	2	7.7% (tows)		
Australia	2019	East Coast Longline			525		12.1% (hooks – SBT Area)	12.2% (no. retained catch)	na
Australia	2019	West Coast Longline			31		12.8%	No observed catch	na
Australia	2020	East Coast Longline			418		11.0% (hooks – SBT Area)	7.7% (no. retained catch)	na
Australia	2020	West Coast Longline			19		12.1%	No observed catch	na

^aAustralian purse seine and towing observer statistics are for the SBT fishing year December–November

^bYear 1 of the 2009–11 season (2009–10); ^cYear 2 of the 2009–11 season (2010–11); na = not available

Appendix F: Number of biological samples taken in observer programs (year and sector)

Country	Year	Sector	Otoliths	Sex	Tags	Stomach contents	Length measurement
Australia	2002	Longline	0	124	165	0	300
Australia	2003	Longline	0	51	229	1	388
Australia	2004	Longline	5	62	0	5	187
Australia	2004–05	Purse seine	2	2	0	0	3
Australia	2005	Longline	63	189	19	12	264
Australia	2005–06	Purse seine	46	46	0	0	23
Australia	2006	Longline	0	4	1	0	32
Australia	2006–07	Purse seine	9	17	0	16	19
Australia	2007	Longline	9	41	0	0	42
Australia	2007–08	Purse seine	4	4	0	0	4
Australia	2008	Longline	0	84	0	1	99
Australia	2008–09	Purse seine	14	14	0	0	14
Australia	2009	Longline	0	746	0	0	810
Australia	2009–11 ^a	Purse seine	3	3	0	0	3
Australia	2010	Longline	0	563	0	0	563
Australia	2009–11 ^b	Purse seine	4	5	0	0	5
Australia	2011	Longline	0	255	0	0	255
Australia	2011–12	Purse seine	5	8	0	0	8
Australia	2012	Longline	0	70	0	0	69
Australia	2012–13	Purse seine	4	6	0	0	119
Australia	2013	Longline	0	1089	0	0	1089
Australia	2013–14	Purse seine	1	1	0	0	1
Australia	2014	Longline	0	290	0	0	290
Australia	2014–15	Purse seine	26	21	0	0	27
Australia	2015–16	Purse seine	3	15	0	0	15
Australia	2016–17	Purse seine	0	23	0	0	23
Australia	2017–18	Purse seine	0	26	0	0	26
Australia	2018–19	Purse seine	0	8	0	0	8
Australia	2019–20	Purse seine	0	0	0	0	1

^aYear 1 of the 2009–11 season (2009–10)

^bYear 2 of the 2009–11 season (2010–11)

Appendix G: An overview of the Australian SBT observer program

Observer sources and training

AFMA has recruited and trained observers since its establishment in 1992. Approximately 15 observers are currently employed in the AFMA observer program. They are sourced from universities and maritime industries from around Australia and require the ability to live and work at sea, have demonstrated experience in collecting biological data at sea, and have experience in fisheries research methodologies and collection of associated scientific data. Observers also hold marine radio operators certificate of proficiency (or similar qualifications and/or experience), a sea safety certificate and medical certificate, and have completed an AFMA observer training course.

In 2008 and 2009, in addition to the independent AFMA observers, an international observer from South Africa (Capricorn Fisheries Monitoring) was deployed. No international observers have been used since then.

AFMA implemented a trial of e-monitoring (i.e. on-board, fixed-mount digital video cameras) in the Eastern Tuna and Billfish Fishery in 2009–10 to evaluate the effectiveness of this technology for a range of fishery monitoring purposes and to conduct a cost–benefit analysis (Piasente et al. 2012). E-monitoring became compulsory for the Australian longline fleet from 1 July 2015. E-monitoring replaces human observers for all in-zone observer requirements. At least 10% of video footage of all hauls will be reviewed to verify the accuracy of logbooks which are required to be completed for 100% of shots. This review rate can be increased to target specific events or behaviours.

Purse seine fishery – Great Australian Bight 2019–20

The purse seine observer program for the 2019–20 Australian SBT fishing season monitored fishing and tow operations in 36°16'–38°00'S and 135°36'–139°10'E in January, February and March 2020. Two Australian observers monitored 14 purse seine sets where fish were retained. Fish were released from one shot as they were deemed to be too small. This represents 9.9% of the total sets in which fish were taken in 2019–20. From these observations an estimated 437 t of SBT were caught during observed sets, representing 10.3% of the estimated tonnage caught. An estimated 36 t of fish were released alive because the fish were too small. There were no mortalities observed during purse seine operations. Two observers monitored two tow operations and recorded one SBT mortality during the towing operation, which was sampled.

Longline fishery

In 2020, in the ETBF, south of 30°S and during the months of May to September, e-monitoring monitored 97,450 hooks of a total of 889,711, representing 11.0% observer coverage of longline effort during the SBT migration. For the fishery as a whole, 9.8% of hooks were observed in 2020. The observed total catch number of SBT was 1,086 individuals, of which 1,044 were retained, 42 fish were discarded (19 of which were reported as released alive, 9 were dead or damaged and 14 of which were reports as having an undetermined life status) and none were

tagged. ETBF logbooks for 2020 showed 13,633 SBT (688.8 t) were retained and 1,615 (10.6%) were released.

During 2020, there was 12.1% observer coverage of longline hook effort in the WTBF. Two longline vessels participated in this fishery in 2020.

Scientific observer program design and coverage

The target coverage for the SBT purse seine fleet operating out of Port Lincoln is 10% of the total catch and effort for the fishery. Observers monitor 100% of all fishing operations while on board. Most of the Australian SBT purse seine effort has historically taken place in an area between 33–35°S and 131–133°E, although this has changed slightly in recent years (see Figure 6).

The observers in the purse seine fishery in the 2019–20 fishing season spent 53 days at sea, and observed purse seiner activities for 10 days and tow activities for 21 days. The remainder of the days were spent in transit, searching or lost due to rough weather.

Observer data collected

Effort data

In 2020, two observers monitored 14 purse seine sets where fish were retained. Fish were released from one shot as they were deemed to be too small. Fishing operations observed in the purse seine sector were based in the Great Australian Bight between 36°16'–38°00'S and 135°36'–139°10'E (see Figure 6). The observed sets where fish were retained represent 9.9% of all sets in the fishery where fish were retained for the 2019–20 season.

Data were gathered on vessel characteristics, fishing gear and equipment. Comprehensive operational and environmental information were recorded for each set that occurred while the observer was on board. This included information on searching, chumming, setting and hauling activities. Information on chumming operations by the fishing vessel and associated chumming vessels was also recorded. In addition, observers recorded information on the movement of some spotter aircraft and their time in the area preceding sets.

Two tows were observed and data collected on the number of SBT mortalities and the date they occurred. Data were also collected on:

- Towing methods
- Average towing speed
- Cage number and diameter
- Maximum cage depth
- Average weight of SBT transferred
- Estimated number of SBT
- Methods of counting and verifying fish counts.

Australia's 2019–20 Southern Bluefin Tuna Fishing Season




Figure 6 The number of sets recorded in the SBT purse seine fishery from December 2019 to April 2020. The hatching represents areas where observations took place.

Catch data

Observers recorded catch composition and fate of target and bycatch species where possible during all observed sets. The time at start and end of observation, the observed catch in estimated number and estimated weight for SBT and all other species were recorded where possible.

Because fish are taken alive for farming purposes in the purse seine sector, it is not possible to obtain actual weight or length information at the time of catching the SBT. Consequently, both catch data and observed catch data are estimates only and these are reported below.

In total, an estimated 437 t of SBT were caught during observed sets. This observed catch accounted for 10.3% of the total estimated catch of 4,224 t. The actual tonnage caught by purse seine vessels in the 2019–20 fishing season was 4,586 t.

The total number of other significant species observed caught and retained by longline, south of 30°S and during the months of May to September, were albacore (1,953), yellowfin tuna (1,108), Ray's bream (154), bigeye tuna (152) and swordfish (109). No other species were observed caught in the purse-seine fishery.

Length frequency data

It is not possible for onboard observers to obtain length measurements for the live SBT catch, but observers are required to take biological samples from all SBT mortalities. During purse seine and towing operations, the observers recorded one SBT mortality, which was sampled. The fork length for this fish was 97 cm.

Biological data

No otoliths were obtained from the seven sampled mortalities in the purse operations. No otoliths were obtained from the one sampled mortality in the tow operation. There is also an ongoing project to collect otoliths from farm mortalities.

A weight was not obtained for the single fish that was sampled during purse seine and towing operations.

Tag return monitoring

There were no tagged SBT observed in 2020.

Appendix H: Australian southern bluefin tuna research projects

The projects below represent the major research investment areas by Australia for the past several years that directly contribute to the work of the Commission.

Category	Project title	Year(s)	Amount (AU\$)
Data supporting	Tagging juvenile SBT off South Africa	2006	\$100,000
SBTF	CCSBT conventional tagging programme	2006	\$170,832
	Archiving of hard parts for SBT in 2006/07	2006	\$28,231
	Archiving of hard parts for SBT in 2007/08–2009/10	2007	\$307,302
	Archiving of hard parts for routine ageing and developing age-length keys for the Australian SBT surface fishery 2010/11–2012/13	2010	\$304,643
	Development of SBT catch and effort monitoring program	2009	\$150,000
	Aerial survey in the Great Australian Bight (GAB) 2008	2008	\$427,274
	Aerial survey in the Great Australian Bight (GAB) 2009	2009	\$582,440
	Aerial survey in the Great Australian Bight (GAB) 2010	2010	\$621,625
	Aerial survey in the Great Australian Bight (GAB) 2011	2011	\$753,208
	Aerial survey in the Great Australian Bight (GAB) 2012	2012	\$778,120
	Aerial survey in the Great Australian Bight (GAB) 2013	2013	\$808,360
	Aerial survey in the Great Australian Bight (GAB) 2014	2014	\$661,635
SBT Research	Acoustic monitoring of juvenile SBT in the GAB	2002–07	\$150,000
	Spatial interactions among juvenile SBT at a global scale: a large-scale archival tag experiment	2003–11	\$2,549,000
	Analysis of overcatch data	2006	\$108,553
	Monitoring of the Japanese SBT market	2006	\$70,000
	Fishery-independent estimate of spawning biomass of southern bluefin tuna through identification of close-kin using genetic markers	2006–11	\$1,491,146
	Australian farm research program	2007	\$200,000

Australia's 2019–20 Southern Bluefin Tuna Fishing Season

	Assessing operational feasibility of stereo video and evaluating monitoring options for the SBT farm sector	2007	\$395,130
	Further monitoring of the Japanese SBT market	2008	\$220,000
	Tasman residency and spawning migrations of adult SBT	2008–09	\$165,000
	Management Procedure development	2009	\$224,899
	SBT stereo-video project	2010	\$75,000
	Developing a management procedure-based recovery plan for SBT	2011	\$358,130
	Otolith chemistry of juvenile southern bluefin tuna	2011–12	\$20,000
	Investigating new data sources and spatial operating models	2011–13	\$315,000
	Automation of measurements from underwater stereo-video	2011–14	\$30,000
	Forecasting spatial distribution of SBT habitat in the GAB (proposal)	2013–14	\$145,948
	Updating the CCSBT operating model and intersessional science	2014–15	\$529,301
	Intersessional science – Impact of no 2015 aerial survey on the MP	2015	\$118,385
	Intersessional science – Updated stock assessment and MP work	2017	\$180,000
	Intersessional science – New MP work	2018	\$180,000
	Intersessional science – MP development	2019	\$185,000
	Intersessional science – MP development and stock assessment	2019–21	\$360,000
Capacity building	Monitoring of longline catch of SBT landed in Indonesia	2003–04	\$130,508
	Monitoring of longline catch of SBT landed in Indonesia	2004–05	\$112,628
	Monitoring of longline catch of SBT landed in Indonesia	2006–08	\$397,389
Recreational fishing	Quantifying the recreational catch of SBT off the Victorian coast	2010–11	\$212,000
	Developing a methodology for obtaining regular, statistically robust estimates of recreational and charter fishing catch	2011–12	\$100,000
	Development of methods for obtaining national estimates of recreational catch	2013–15	\$570,000
	Capture-induced physiological stress and post-release survival of recreationally caught Southern Bluefin Tuna	2013–15	\$383,486
	National survey for recreational catch of southern bluefin tuna	2018–20	\$2,311,936

References

- Anonymous 2011, *Technical assessment of the 2011 commercial trial of stereo-video in the Australian southern bluefin tuna farm sector*, Working paper CCSBT-CC/1110/11, Sixth meeting of the Compliance Committee of the Commission for the Conservation of Southern Bluefin Tuna, Bali, Indonesia, October 2011
- Caton, AE & Ward, PJ 1996, 'Arrangements on fisheries between the governments of Australia and Japan on tuna longlining, Annex to Ward PJ (Ed) (1996) Japanese longlining in eastern Australian waters 1962–1990', Bureau of Resource Sciences, Canberra
- Caton, AE, Ward, PJ, Colgan, MK, Williams, KF, Ramirez, C & Skousen, T 1995, *The Australian 1989–90 to 1994–95 southern bluefin tuna seasons*, Working Paper SBFWS/95/1, first meeting of the Scientific Committee of the Commission for the Conservation of Southern Bluefin Tuna, Shimizu, Japan, July 1995
- Forbes, E, Tracey, S & Lyle, J 2009 'Assessment of the 2008 recreational gamefish fishery of southeast Tasmania, with particular reference to southern bluefin tuna', Tasmanian Aquaculture and Fisheries Institute, University of Tasmania, Hobart
- Moore, A, Hall, K, Khageswor, G, Tracey, S, Hansen, S, Ward, P, Stobutzki, I, Andrews, J, Nicol, S & Brown, P 2015, 'Developing robust and cost-effective methods for estimating the national recreational catch of southern bluefin tuna in Australia', FRDC report 2012/022.20, ABARES, Canberra
- Piasente, M, Stanley, B, Timmiss, T, McElderry, H, Pria, M, & Dyas, M 2012, *Electronic onboard monitoring pilot project for the Eastern Tuna and Billfish Fishery*, FRDC Project 2009/048, Australian Fisheries Management Authority, Canberra
- Shafait, F, Harvey, ES, Shortis, MR, Mian, A, Ravanbakhsh, M, Seager, JW, Culverhouse, PF, Cline, DE & Edgington, DR 2017, Towards automating underwater measurement of fish length: a comparison of semi-automatic and manual stereo-video measurements, ICES Journal of Marine Science, vol. 74, pp. 1690–1701
- Tracey, S, Hartmann, K, McAllister, J, Conron, S & Leef, M 2016 *Capture-induced physiological stress and post-release survival of recreationally caught Southern Bluefin Tuna*, Final report FRDC project 2013-025, Institute for Marine and Antarctic Studies, Hobart, Australia.
- Tracey, S, Lyle, JM, Ewing, G, Hartmann, K & Mapleston, A 2013, *Offshore recreational fishing in Tasmania 2011/12*, Institute of Marine and Antarctic Studies, University of Tasmania, Hobart.
- Tracey, SR, Lyle, JM, Stark, K, Gray, S, Moore, A, Twiname, S & Wotherspoon, S 2020, *National survey of recreational fishing for southern bluefin tuna in Australia 2018–19*, University of Tasmania, Hobart