



Analyses of tag return data from the CCSBT SRP tagging program - 2008

**Paige Eveson
Tom Polacheck**

**Prepared for the CCSBT 5th Management Procedure Workshop 2-7 September and the
13th Meeting of the Extended Scientific Committee 8-12 September 2008
Rotorua, New Zealand**

Table of Contents

Abstract.....	2
Introduction.....	3
Methods.....	3
Data.....	4
Estimation Model.....	5
Reporting Rates.....	7
Tag Shedding.....	7
Natural Mortality Rates.....	8
Results and Discussion.....	8
Location of Longline Returns.....	8
Fishing Mortality Rate Estimates.....	9
Returns from the First Season for December Releases.....	10
Summary.....	10
Acknowledgements.....	12
Literature Cited.....	13

List of Tables

Table 1: The set of alternative reporting rate estimates considered for the surface fishery, taken from Table 4 of Hearn et al. (2008).	14
Table 2: Shedding rate estimates by individual taggers and by tagger groups (results provided by Dr. W.S. Hearn, CSIRO Marine and Atmospheric Research).....	14
Table 3: Age-specific natural mortality rates used in the estimation of fishing mortality rates.	14
Table 4: Total number of tag releases and reported recaptures (i.e. returns) by cohort (includes returns to date from the 2008 fishing season).	15
Table 5: The number of releases by age and corresponding returns by age for the 1998-2005 cohorts. Note that rows with less than 20 releases have not been included.....	15
Table 6: Percent of (reported) longline recaptures that occurred in the Tasman Sea (defined as east of 142°E). Note that the RMP results have not been updated for returns in 2005/2006 and beyond.	16
Table 7: Percent of (reported) longline recaptures from the Tasman Sea (defined as east of 142°E) that occurred in the Australian Fishing Zone (AFZ) for the 1990s RMP tag releases.	16
Table 8: Estimates of age-specific fishing mortality rates (F) for different cohorts derived from tags released in the waters around southern Australia. Results are presented separately for tags released at different ages. Lower and upper 5% refer to the lower and upper 5th percentiles from the bootstrap estimates (i.e. the 90% confidence interval). Results were derived using reporting rate vector 1 from Table 1 for the surface fishery, group shedding rates from Table 2, natural mortality vector 1 from Table 3, and a reporting rate of 0.65 for the longline fisheries. (F estimates > 0.4 are shaded.)	17
Table 9: Comparison of age-specific fishing mortality rate estimates for a range of scenarios, as described below. (F estimates > 0.4 are shaded.)	19
Table 10: First year recaptures of fish released in December in the Great Australian Bight by age of release and fishing year. Estimated percent caught is based on weighted reporting rate estimates in Table 1, with no allowance for tag shedding.	21

List of Figures

Figure 1: Comparison of fishing mortality rate (F) estimates at ages 3 and 4 obtained from the SRP tagging with those obtained from the 1990s RMP tagging. Estimates for the 1990s tagging were obtained using a Brownie model that integrates data from all release ages.....	22
--	----

Abstract

An updated summary of the release and recapture data from the CCSBT SRP tagging program is presented, along with some updated estimates of fishing mortality rates. A tag attrition model was again used to estimate cohort and age-specific fishing mortality rates for different groups of tag releases conditional on estimates of natural mortality, tag shedding and reporting rates (the latter three derived from separate analyses).

The results continue to show very high estimates of fishing mortality rates for ages 3 and above (many over 0.5) over the past four years based on fish tagged at age 2 and older; although the age 3 estimate for 2007 is somewhat lower (0.26). These results hold true for a range of scenarios using alternative reporting rate estimates for the surface and longline fisheries and alternative natural mortality rate vectors. Comparison of these results with those from the 1990s RMP tagging indicates that the fishing mortality being experienced by tagged fish has substantially increased. Even the age 3 estimate for 2007 remains above levels experienced in the 1990s.

There continues to be a marked lack of returns, and thus lower estimates of fishing mortality, from fish tagged at age 1 compared to those tagged at ages 2 and above. This phenomenon was not observed in the tag returns from the 1990s releases, and suggests that one year old fish found in WA (where the majority of age 1 fish are tagged) are no longer entering the GAB in substantial numbers. These same one year old fish do not appear to be entering the longline fisheries either. Furthermore, the spatial distribution of longline returns from the 2000s tagging consistently show a much smaller percentage of tagged fish moving into the Tasman Sea in recent years than in the 1990s. These spatial changes have now been observed in five years of release and recovery data, suggesting that they are not simply outliers.

Changes in the exploitation rate estimates and spatial patterns of returns between the 1990s and 2000s suggest possible negative consequences in terms of current stock status (e.g. increased exploitation rates and possible range contraction). Only through continued and improved tagging experiments can the longer term consistency, implications, and underlying sources of these observed changes be understood. As such, it would seem critical that the SRP tagging program that was suspended last year be resumed and improved.

Introduction

As part of the Scientific Research Program (SRP), the CCSBT initiated a large scale tagging program to estimate juvenile fishing mortality rates beginning in 2000/2001 fishing season. The basic design of the tagging program was similar to that conducted in the 1990s as part of the CSIRO/NRIFS Recruitment Monitoring Program with the aim to tag multiple cohorts at different ages in several years. This paper provides an updated analysis of the data collected to date in the SRP tagging program, including some estimates of fishing mortality rates obtained using the same approach as in Polacheck and Eveson (2005, 2006, 2007).

Methods

Much of the Methods section remains unchanged from last year's report (Polacheck and Eveson 2007), but it is repeated here in full for easy reference.

Data

Tagging in the first year was only done off Western Australia (WA) with 1 and 2 year old fish being tagged. In all subsequent years, tagging was conducted in both WA and South Australia (SA) with almost all of the fish tagged being between ages 1 and 3 (i.e. less than 2% of the fish tagged are estimated to be older than age 3).

Some of the release and return data are considered unreliable for estimating mortality rates; therefore we applied the following screening process to the data prior to analysis.

For the release data:

- Only fish released into the wild were included (i.e., we excluded data from fish that were released into farms as part of a tag seeding program).
- Only releases where the fish was caught by pole and line were included. This method of catching fish is least likely to cause lasting injury to the fish.
- Only releases for which both tags were recorded as being inserted correctly were included to reduce the chance of tag shedding biasing our analyses.
- Only fish for which the injury due to tagging was regarded as slight were included to reduce the chance of fish mortality due to tagging biasing our analyses.
- Only fish whose length was recorded at the time of tagging were included because our analysis uses age of release, which is estimated based on length.

For the recapture data:

- Only recaptures corresponding to releases that met the above release criteria were included.
- Only recapture records from fish caught in the wild were included. For tagged fish that are harvested from the farms, the database has two records: one corresponding to the original capture from the wild and one corresponding to the harvest from the farm. For the purposes of estimating fishing mortality we are only interested in the information (date and location) for the capture from the wild.

A fish's age at tagging was estimated from its length using cohort slicing and the growth curve currently adopted by the CCSBT (Anon. 2001). SBT grow rapidly as juveniles so there is good separation between length distributions at the ages being tagged, and therefore the number of aging errors should be small. All tagging was done between December and April, so the release ages were adjusted in order that fish tagged in December from a given year-class/cohort were assigned the same age as those tagged after December. The recapture age was calculated using the age of release and the time between release and recapture.

Recapture ages were also adjusted so that fish from a given cohort caught in November or December were given the same age as those caught after December.

As discussed below, results from separate analyses of tag shedding rates performed by Dr. W.S. Hearn (CSIRO Marine and Atmospheric Research) are used in the estimation of mortality rates. In addition to the above data screening, Dr. Hearn excluded tag returns if the recapture year or month within year was uncertain, or if the day within month was uncertain for recaptures at liberty less than 270 days. Also, data sets associated with a tagger were only

analysed if there were 30 or more acceptable recaptures in the set. Data associated with the remaining taggers were pooled into a set we call “tagger” ZZ.

The data used in this paper were taken from the most recent version of the CCSBT tag database (2008-05-26). Recaptures from the 2007/2008 fishing season (i.e., recaptures since November 2007) have been excluded since the data for this year are still very incomplete (only a small fraction of fish had been harvested from the farms at the time the data were compiled for analysis).

Estimation Model

A basic tag attrition model was used to estimate cohort and age-specific fishing mortality rates for different groups of tag releases. This model was chosen because it provides a direct estimate of the fishing mortality rate for those fish tagged independent of any assumptions about mixing. This is seen as a first step to evaluate the consistency of estimates from different releases prior to developing a more integrated estimation model (e.g. a Brownie model).

We define two seasons: season 1 runs from January 1 to June 30 and corresponds to the Australian surface fishery; season 2 runs from July 1 to December 31 and corresponds to the longline fishery. For convenience, the model assumes all releases occurred in season 1 on January 1. In addition, the model follows the convention used in the CCSBT Management Procedure operating model in which all fishing is assumed to occur either on January 1 (season 1) or July 1 (season 2). All returns from the Australian surface fishery were assumed to occur in season 1 and all longline returns were assumed to occur in season 2. Natural mortality is assumed to occur at a constant rate throughout the year (i.e., it is evenly split between the two seasons since they are of equal length).

Because there are two seasons per year, it is convenient to work in terms of time periods taking values $t = 1, 2, 3, \dots$, where season 1 corresponds to odd time periods, season 2 corresponds to even time periods, and a year consists of a consecutive odd and even time period.

Let

$$N_{c,a,g,t+1} = (N_{c,a,g,t} - \widehat{R}_{c,a,g,t})e^{-0.5m_{a^*}} \quad (1)$$

where

$$\begin{aligned} N_{c,a,g,t} &= \text{the number of tagged fish alive at the start of time period } t \text{ from} \\ &\quad \text{fish tagged from cohort } c \text{ at age } a \text{ by tagger group } g; \\ \widehat{R}_{c,a,g,t} &= \text{the estimated number of tagged fish caught in time period } t \text{ from} \\ &\quad \text{fish tagged from cohort } c \text{ at age } a \text{ by tagger group } g; \\ m_{a^*} &= \text{natural mortality for fish of age } a^* = y - c, \text{ where } y \text{ denotes the} \\ &\quad \text{year corresponding to time period } t. \end{aligned}$$

The subscript g in the above equation is necessary if one wishes to examine results for different groups of taggers. This could be all taggers, a group of a few taggers, or a single tagger. In the current paper, we only present results using all taggers, but in past we have compared results for different groups of taggers (e.g. Polacheck and Eveson 2007).

The number of recaptured tagged fish, $\widehat{R}_{c,a,g,t}$, in equation 1 is not simply the number of tags actually returned but is estimated to take into account both tag shedding and non-reported tags. Specifically, $\widehat{R}_{c,a,g,t}$ is estimated by

$$\widehat{R}_{c,a,g,t} = \frac{\sum_{k \in g} (R_{c,a,k,t} / \gamma_{c,a,k,t})}{\lambda_t}$$

where

$R_{c,a,k,t}$ = the actual number of reported tag returns in time period t from fish tagged from cohort c at age a by a tagger in sub-group k of tagger group g ;

$\gamma_{c,a,k,t}$ = the probability that a fish tagged from cohort c at age a by a tagger in sub-group k has at least one tag still attached at the beginning of time period t ;

λ_t = the tag reporting rate in time period t .

Recall that all fish have been double-tagged. The probability of a tagged fish still having at least one tag attached at the time of capture,

$\gamma_{c,a,k,t}$, is given by

$$\gamma_{c,a,k,t} = 1 - [1 - Q_{a,k}(\tau)]^2$$

where

$Q_{a,k}(\tau)$ = the probability of a tag still being attached to a fish tagged at age a by a tagger in sub-group k after the fish has been at liberty for time τ . Note that τ is a function of c and a (which together define the time period of release) and t (the time period of recapture).

Finally, an estimate of the annual fishing mortality rate in year y , corresponding to time periods t and $t+1$ (where t is odd), for fish from cohort c (i.e. age $a^* = y - c$) can be calculated from the ratio of the estimated number of tagged fish alive at the start of year $y+1$ (time period $t+2$) to the estimated number of tagged fish alive at the start of year y (time period t). A separate value can be calculated corresponding to fish tagged at age a by a tagger in tagger group g . Thus,

$$f_{c,a,g,y} = - \left[m_{a^*} + \log \left(N_{c,a,g,t+2} / N_{c,a,g,t} \right) \right]$$

Bootstrap confidence intervals for $f_{c,a,g,y}$ were calculated by sampling the releases at age a from cohort c by tagger group g along with the associated recapture data with replacement and calculating the estimates of $f_{c,a,g,y}$ for each bootstrap sample. The confidence intervals presented are based on 1000 bootstrap replicates and treat each tag release as independent. This may underestimate the actual uncertainty if releases from the same school tend to stay together. The bootstrap estimates are also conditional on the estimates of reporting rates, shedding rates and natural mortality rates.

Reporting Rates

Estimates of the reporting rate in the Australian surface fishery are available for the 2002/03 to the 2007/08 fishing season from tag seeding experiments conducted in these years (Polacheck and Stanley 2004, 2005; Polacheck et al. 2006a; Hearn et al. 2007; Hearn et al. 2008). The estimates declined steadily from 0.64 in 2002/03 to 0.30 in 2007/08, then increased slightly in 2007/08 to 0.43 (Table 1, vector A1). The low estimates in recent years can yield unrealistically high values for some fishing mortality rates, and has raised the question of whether in fact the tag seeding results are providing unbiased estimates of the reporting rate (Polacheck and Eveson 2007). Hearn et al. (2008) further discuss this issue and suggest a number of alternative reporting rate vectors for consideration (Table 1, vectors A2-A5). Brief descriptions of the alternative reporting rate vectors are as follows:

- A1. “Best estimates” from the tag seeding results.
- A2. Assumes that the reporting rate in 2002/03 was 1.0 and that the difference between the estimate of 0.64 from the tag seeding was due to high initial shedding of both tags (i.e. a lack of independence in shedding) associated with tagged fish being in cages. Further assumes that the rate of high initial shedding is constant across years and re-adjusts the other reporting rates accordingly.
- A3. Estimates based only on Tagger 4; the most consistently used tagger and also one with extensive experience.
- A4. Assumes reporting rates have been constant and uses the rate of return from the re-release of wild tagged fish from the 40 fish samples as an estimate of the reporting rate (see Polacheck and Eveson 2007 for details).
- A5. Assumes reporting rates were the same in 2002/03 to 2003/04 and 2004/05 to 2006/07 and uses the rate of return from the re-release of wild tagged fish from the 40 fish samples for these two periods to estimate the reporting rates.

Insufficient information was available to estimate reporting rates from the longline fisheries. Estimates of reporting rates from longliners were substantially below those in the surface fishery in the 1990s. Reporting rates for Japanese longliners in the 1990s ranged from 0.07 to 0.49 (Eveson and Polacheck 2005). There were no data to directly estimate reporting rates for Taiwanese vessels. In the absence of any direct data, results were explored for two values, namely 0.65 and 0.30, to provide an indication of the sensitivity of the results to the value assumed. The same value was used for all ages and years. Note, however, that unless the reporting rates were the same in the different longline fleets, the reporting rate would in fact vary with age and year even if the reporting rate was constant over time within a fleet; this is because the proportion of the total longline catch of a given age class by a given fleet varies among years (Pollock et al. 2002).

Tag Shedding

Tag shedding rates provided by Dr. Hearn were based on an analysis of the tag shedding data (number of recaptures with one tag versus two tags still attached) for taggers who participated in the SRP tagging program. Dr. Hearn applied the method of Kirkwood and Walker (1984) to estimate shedding parameters. The retention function (i.e., the probability of a tag still being attached after being at liberty for time τ) was assumed to have the form

$$Q_{a,k}(\tau) = \xi_{a,k} \exp(-\Omega_{a,k} \tau)$$

where $\zeta_{a,k}$ is the fraction of tags immediately retained (i.e. $1 - \zeta_{a,k}$ are immediately shed) for fish tagged at age a by a tagger in sub-group k , and $\Omega_{a,k}$ is the continuous shedding rate. The model allows for tag shedding to vary between tagger groups (which may be individual taggers) and between fish released at different ages. The retention function was assumed to be the same for both tags on a given fish. Table 2 provides the estimates of the parameters for this retention function when fitted to the SRP tag return data. This table provides estimates for individual taggers as well as for groups of taggers which do not differ significantly in their tag shedding parameters. Only the estimates for the groups of taggers are used in the estimates of fishing mortality rates presented here, but the results are very similar if estimates for individual taggers are used. In the notation above, each set of taggers constitutes a potential sub-group k .

In Polacheck and Eveson (2005), we also considered the potential effect of age-specific shedding. There were only sufficient data to meaningfully perform these calculations for two taggers. Only for one of these was the difference significant and the differences had only a minor effect on the overall results. As such, we have not subsequently updated these estimates.

Natural Mortality Rates

Two age-specific natural mortality rate vectors were used in the calculation of the fishing mortality rates (Table 3) to provide a measure of the sensitivity of the estimates to assumptions about natural mortality. These are two of the vectors being used in the conditioning and projections being undertaken with the SBT Management Procedure operating model.

Results and Discussion

Table 4 provides an updated summary of the number of releases and recaptures by cohort. Note that it was decided at the 2007 CCSBT SAG and SC meetings to suspend the conventional tagging program for at least one year; thus, only a small number of juvenile SBT (~400) were tagged in the 2008 season as part of field exercises for other research programs. The low number of recaptures from the 2005 cohort and beyond reflect the fact that it is still too early to expect any substantial numbers of returns from these releases. Given the current fisheries, only significant numbers of recaptures are expected from fish ages 3 and older. Since most of the returns from this year's Australian surface fishery are not yet available, the tagging data are not yet informative even for releases from the 2005 cohort. As such, the focus of the results presented are for the 1999-2004 cohorts (the number of releases for the 1998 cohort are too small to provide meaningful results).

Table 5 provides a breakdown of the release and recapture data by cohort, age at release and age at recapture. As noted in previous years, the percent of returns from fish released at age 1 tends to be very low compared to fish released at ages 2 and 3 from the same cohort. This was not observed in the 1990s RMP tag returns, but continues to be a persistent feature of the SRP tag returns; possible reasons were explored and discussed in Polacheck and Eveson (2007).

Location of Longline Returns

It has become evident in recent years that the proportion of longline tag returns coming from the Tasman Sea has been much lower in the 2000s compared to the 1990s (Table 6). As

discussed in Polacheck and Eveson (2007), this in part reflects changes in the spatial distribution of fishing effort within the Tasman Sea. In particular, a substantial fraction of longline fishing effort occurred in the Australian Fishing Zone (AFZ) in the 1990s as a result of joint venture operations and bi-lateral access arrangements that allowed vessels to fish within the AFZ (Table 7). These arrangements ceased in 1998 and thus there has been little recent fishing effort in the areas where substantial numbers of the 1990s returns came from. However, it seems unlikely that these factors are sufficient to explain the large differences in the spatial distribution of longline returns in the 1990s compared to the 2000s (refer to Polacheck and Eveson 2007 for a detailed discussion).

Fishing Mortality Rate Estimates

Sufficient release and return data (e.g. at least ~400 releases at a particular age and at least one year of full recoveries) exist to derive age-specific fishing mortality rate (F) estimates for cohorts 1999 to 2004 tagged at ages 1 to 3 (with the exception of cohort 1999). Table 8 contains the age-specific F estimates, along with bootstrapped 90% confidence intervals, obtained using reporting rate vector A1 from Table 1 for the surface fishery, group shedding rate parameters from Table 2, natural mortality vector 1 from Table 3, and a reporting rate of 0.65 for the longline fisheries. It should be noted that estimates of fishing mortality rates based on returns from the same year of release (e.g. the F estimates for age 2 based on age 2 releases) can be highly misleading in terms of being representative of the fishing mortality experienced by a cohort because the releases may have occurred before, during or after the main period of fishing, and the distribution of releases would also affect the number of returns. However, they do provide a measure of the fishing mortality rate experienced by the set of tagged fish and in this sense can still be informative.

Some high F estimates are evident in Table 8 (values greater than 0.4 are shaded). In particular, fishing mortality rates at ages 3 and above tend to be high in years 2004 to 2006 for fish tagged at age 2 and older (e.g., ranging from 0.518 to 1.046, with a mean of 0.77, for age 3 fish tagged at age 2; and ranging from 0.282 to 1.140, with a mean of 0.70, for age 4 fish tagged at ages 2 and 3). In a few cases, the estimates seem unrealistically high (e.g., greater than 1.0). The most recent results, i.e., the estimates for 2007, suggest a lower fishing mortality rate for age 3 fish (0.26), but an extremely (and unrealistically) high mortality rate for age 4 fish (2.1).

Also evident is that the F estimates for ages 3 and above for fish tagged at age 1 are markedly lower than the 'equivalent' estimates from fish tagged at older ages (i.e., F estimates for a given recapture year and age are much lower when estimated using releases at age 1 than when estimated using releases at ages 2 and above.) Importantly, no such differences were observed in the F estimates from the 1990s tagging data. This anomaly has been noted in past and was discussed in detail in last year's report (Polacheck and Eveson 2007). As such, we will not repeat the discussion here except to say that the reason for this inconsistency has not been resolved.

Table 9 provides a comparison of estimates of fishing mortality rates for a range of scenarios. Specifically, scenario 1 contains the estimates from Table 8, as described above. Scenarios 2-4 are the same as scenario 1 except: scenario 2 uses reporting rate vector 4 from Table 1 for the surface fishery; scenario 3 uses a reporting rate of 0.30 for the longline fisheries; and scenario 4 uses natural mortality vector 2 from Table 3. As would be expected, the estimates are somewhat higher with a lower reporting rate for the longline fishery (scenario 3) and with higher natural mortality rates (scenario 4). The estimates for scenario 2 compared to

scenario 1 can vary since reporting rate estimates are higher for A4 than A1 in some years and lower in others. Overall, however, the very high and, in some cases, seemingly “unrealistic” fishing mortality rate estimates are evident for all scenarios.

Figure 1 compares estimates of fishing mortality rates for ages 3 and 4 obtained from the SRP tagging with those obtained from tagging experiments conducted in the 1990s as part of the collaborative CSIRO NIRFSF Recruitment Monitoring Programme (RMP). For the SRP estimates, F estimates are shown for different ages of release (excluding estimates that are for the same age as the age of release). The RMP estimates for the 1990s are based on a Brownie estimation model that integrates data from all release ages, since no substantive inconsistencies were found among the recoveries for different ages at release (see Polacheck et al. 2006 for more discussion on this). While the estimators used for the 1990s and 2000s releases are different, the general tagging locations and methods were the same. As such, Figure 1 provides an overview of the differences between these two periods. Evident in Figure 1 is that the fishing mortality being experienced by the tagged fish is substantially greater in the 2000s than it was in the 1990s. While interpretation of the differences are confounded by the relatively lower return rates from age 1 releases compared to older ages in the SRP tagging noted above, the results suggest that juvenile exploitation rates at ages 3 and 4, at least for fish found within the GAB, have increased substantially and to high levels. Even the F estimates at age 3 for age 1 releases have tended to exceed the F estimates across all release ages in the 1990s. The estimates for the 2001, 2002 and 2003 cohorts based on age 2 releases are very high (i.e. >0.5). This would be consistent with other indicators suggesting that these cohorts were unusually small. The F estimate at age 3 for the 2004 cohort based on these same age 2 releases has decreased, suggesting that this cohort may be larger (or perhaps better mixed with WA fish). Nevertheless, the estimate is still substantially greater than any of the estimates in the 1990s.

Returns from the First Season for December Releases

Tag returns from releases near the beginning of the fishing season in the GAB can provide an indication of localized exploitation, particularly if tagging does not take place in the immediate vicinity of fishing operations. In 2003 to 2007, some of the SRP tagging operations took place in December in the GAB in inshore areas, whereas fishing operations were concentrated near the shelf edge. Over the next four months, tags were returned from 9%-38% of the age 3, 4 and 5 fish (Table 10). These recapture rates suggest high exploitation rates for fish found in the GAB in December, particularly after taking into account the estimates of reporting rates from tagging seeding experiments (A1 of Table 1).

The extent to which these represent global rates depends in part on the proportion of the age 3 to 5 fish that are in the GAB during the summer months.

Perhaps somewhat surprising in these data are the low levels of returns from age 2 fish tagged in the same location and time period (Table 10). Less than 5% of the fish tagged at age 2 were estimated to have been recovered during the fishing season in spite of the fact that there were sizable catches of 2 year olds in these years.

Summary

The updated results from an analysis of the tag-return data from the CCSBT SRP tagging program continue to suggest high fishing mortality rates (many over 0.5) for fish of age 3 and above in the GAB over the past four years; although the age 3 estimate for the most recent year (2007) is somewhat lower (0.26). Comparison of the SRP tagging results with those from the 1990s RMP tagging indicates that the fishing mortality being experienced by tagged

fish has substantially increased. While interpretation of the differences is confounded by the relatively lower return rates from age 1 releases compared to older ages in the SRP tagging, the results suggest that juvenile exploitation rates for age 3 and 4 fish found within the GAB have increased substantially and to high levels. These higher exploitation estimates have been seen consistently for the last five years. Although the age 3 estimate for 2007 is not as high as in the previous few years, it remains above levels experienced in the 1990s.

A number of apparent anomalous features in the fishing mortality rates from the SRP tagging program were discussed in Polacheck and Eveson (2007). One such anomaly, which was noted again in this report, is the marked lack of returns (and thus lower fishing mortality estimates) from fish tagged at age 1 compared to those tagged at ages 2 and above—a phenomenon that was not observed in the tag returns from the 1990s releases. Other anomalies that were not discussed again in this year's report include: i) evidence of significant tagger effects, with tagger 1 consistently yielding somewhat higher fishing mortality estimates; ii) low estimates of fishing mortality rates at age 2, which appear inconsistent with the catch data from the surface fishery; iii) estimated number of tags returned per 1000 fish caught in the surface and longline fisheries also suggest inconsistencies with the catch data. These issues remain relevant and a full discussion can be found in Polacheck and Eveson (2007).

Comparison of the 1990s and 2000s tag returns also indicates substantial and unexplained changes in the spatial dynamics of juvenile fish. One year old fish found in WA are no longer entering the GAB in substantial numbers. These same one year old fish appear not to be entering the longline fisheries. These differences raise questions as to where these fish are going (or whether they are possibly dying at very high rates from natural causes). Furthermore, the spatial distribution of longline returns from the 2000s tagging consistently show a much smaller percentage of tagged fish moving into the Tasman Sea in recent years than in the 1990s. These changes have now been observed in five years of release and recovery data, suggesting that they are not simply outliers. The underlying cause for these differences is unknown. However, they would be consistent with either a density-dependent induced spatial contraction and/or environmentally driven changes in spatial distribution. Both of these, but particularly the first, would have implications for the stock assessment and advice on the effects of possible management actions.

The differences seen in the estimates of fishing mortality rates and the spatial pattern of returns between the 1990s and 2000s demonstrate the value and importance of the tagging experiments for providing insights into the status of the juvenile component of the SBT stock. The recent results raise a large number of unanswered questions as to the mechanism(s) underlying the changes. However, the changes suggest possible negative consequences in terms of current stock status (e.g. increased exploitation rates and possible range contraction). It seems that only through continued and improved tagging experiments that the longer term consistency, implications, and underlying sources of the observed changes can be understood. Cessation of tagging at this point will break the continuity of the time series of data that have been generated to date, and leave unresolved the persistence of the observed changes. It will also eliminate one of the few fishery independent (i.e. non-CPUE based) indicators for the stock assessments. For these reasons, it would seem critical that the SRP tagging program that was suspended last year be resumed and improved.

Acknowledgements

We wish to acknowledge the large number of individuals that have contributed to the CCSBT SRP tagging program, including the taggers, the crews of the tagging vessels, those involved in promotion and tag recovery activities, and those responsible for the data processing and management aspects. A special thanks to Bill Hearn for his analyses and provision of the tag retention parameters used in this paper and useful discussion on the reporting rates. Funding for this work was provided by AFMA, CSIRO's Wealth from Oceans Flagship and DAFF.

Literature Cited

- Anon. 2001. CCSBT Report of the Fifth Meeting of the Scientific Committee. 19-24 March 2001. Tokyo, Japan
- Eveson, J.P. and T. Polacheck. 2005. Updated estimates of tag reporting rates for the 1990s southern bluefin tuna tagging experiments. CCSBT-MPTM/0502/05.
- Kirkwood, G.P. and M.H. Walker. 1984. A new method for estimating tag shedding rates, with application to data for Australian Salmon, *Arripes trutta esper* Whitley. Aust. J. Mar. Freshw. Res. 35: 601-606.
- Hearn, W., T. Polacheck, C. Stanley and M. Rowlands. 2007. Estimates of reporting rate from the Australian surface fishery based on previous tag seeding experiments and tag seeding activities in 2006/2007. CCSBT-ESC/0709/14.
- Hearn, W., T. Polacheck, C. Stanley. 2008. Estimates of reporting rate from the Australian surface fishery based on previous tag seeding experiments and tag seeding activities in 2007/2008. CCSBT-ESC/0809/21.
- Polacheck, T. and P. Eveson. 2005. Initial analyses of tag return data from the CCSBT SRP tagging program. CCSBT-ESC/0509/21.
- Polacheck, T. and P. Eveson. 2006. Analysis of tag return data from the CCSBT SRP tagging program. CCSBT-ESC/0609/15.
- Polacheck, T. and P. Eveson. 2007. Analysis of tag return data from the CCSBT SRP tagging program. CCSBT-ESC/0709/19.
- Polacheck, T. and C. Stanley. 2004. Update on Tag Seeding Activities and Preliminary estimates of reporting rate from the Australian surface fishery based on tag seeding experiments. CCSBT-ESC/0509/15.
- Polacheck, T. and C. Stanley. 2005. Tag Seeding Activities in 2004/2005 and Preliminary estimates of reporting rate from the Australian surface fishery based on previous tag seeding experiments. CCSBT-ESC/0509/20.
- Polacheck, T., W. Hearn, C. Stanley and M. Rowlands. 2006a. Estimates of reporting rate from the Australian surface fishery based on previous tag seeding experiments and tag seeding activities in 2005/2006. CCSBT-ESC/0609/14.
- Polacheck, T., Eveson, J.P., and Laslett, G.M. 2006b. Estimation of mortality rates from tagging data for pelagic fisheries: analysis and experimental design. Final Report to the Fisheries Research and Development Corporation (FRDC), Deakin West, Australia. FRDC Proj. 2002/015. ISBN 1 921061 03 0.
- Pollock, K., W.S. Hearn and T. Polacheck. 2002. A general model for tagging on multiple component fisheries: an integration of age-dependent reporting rates and mortality estimation. J. Envir. and Ecol. Stat. 9: 57-69.

Table 1: The set of alternative reporting rate estimates considered for the surface fishery, taken from Table 4 of Hearn et al. (2008).

Reporting rate vector	2002/2003	2003/2004	2004/2005	2005/2006	2006/2007
A1	0.64	0.50	0.40	0.30	0.43
A2	1.00	0.78	0.63	0.47	0.67
A3	-	0.63	0.34	0.50	0.47
A4	0.45	0.45	0.45	0.45	0.45
A5	0.67	0.67	0.37	0.37	0.37

Table 2: Shedding rate estimates by individual taggers and by tagger groups (results provided by Dr. W.S. Hearn, CSIRO Marine and Atmospheric Research).

Tagger ID	Initial retention fraction (ξ)	Continuous shedding rate (Ω)	Recaptures with 2 tags	Recaptures with 1 tag	Total number recaptures	
2	0.973	0.102	1872	781	2653	
4	0.902	0.197	841	824	1665	
418	1.000	0.172	48	49	97	
419	1.000	0.279	78	106	184	
444	1.000	0.198	52	45	97	
570	0.895	0.170	97	79	176	
1439	0.781	0.072	261	224	485	
1525	0.891	0.000	175	43	218	
1646	0.892	0.154	231	184	415	
ZZ	0.438	0.000	7	18	25	
Tagger Group						
1	1525	0.891	0.000	175	43	218
2	2	0.973	0.102	1872	781	2653
3	418+444+1439+1646	0.816	0.095	592	502	1094
4	4+419+570	0.908	0.202	1016	1009	2025
5	ZZ	0.438	0.000	7	18	25

Table 3: Age-specific natural mortality rates used in the estimation of fishing mortality rates.

Vector	Age				
	1	2	3	4	5
1	0.3401	0.3028	0.2700	0.2420	0.2153
2	0.4202	0.3703	0.3278	0.2894	0.2538

Table 4: Total number of tag releases and reported recaptures (i.e. returns) by cohort (includes returns to date from the 2008 fishing season).

Cohort	Number releases	Number returns	Percent
1998	50	7	14.0
1999	1190	143	12.0
2000	5789	901	15.6
2001	9899	2095	21.2
2002	10307	1571	15.2
2003	14481	2686	18.5
2004	15154	782	5.2
2005	17897	158	0.9
2006	4068	0	0.0
2007	14	0	0.0

Table 5: The number of releases by age and corresponding returns by age for the 1998-2005 cohorts. Note that rows with less than 20 releases have not been included.

Cohort	Age at release	No. releases	No. returns by age								Total returns	Percent returns
			1	2	3	4	5	6	7	8		
1998	5	44					3	1	1		5	11.4
1999	2	750			11	51	10	3	2	1	78	10.4
	3	23				1		2			3	13.0
	4	414				34	16	8	3	1	62	15.0
2000	1	1921		4	88	19	4	1			116	6.0
	2	492		1	50	37	14	1	2		105	21.3
	3	3276			297	256	68	20	4	1	646	19.7
	4	32				7	5	1	1		14	43.8
	5	68					12	6	2		20	29.4
2001	1	2748		9	129	19	7	3			167	6.1
	2	5869		31	1093	284	80	6	3		1497	25.5
	3	1146			253	103	42	6	2		406	35.4
	4	135				12	10	2			24	17.8
2002	1	3316	1	26	69	25	4				125	3.8
	2	6256		90	704	356	122	3			1275	20.4
	3	720			54	91	23	1			169	23.5
2003	1	2662		33	154	69					256	9.6
	2	8692		102	1222	570	6				1900	21.9
	3	3127			243	285	2				530	16.9
2004	1	7084	1	31	127	3					162	2.3
	2	7591		68	515	4					587	7.7
	3	479			32	1					33	6.9
2005	1	9196	3	33	2						38	0.4
	2	8699		113	7						120	1.4
2006	1	4038										0.0
	2	30										0.0

Table 6: Percent of (reported) longline recaptures that occurred in the Tasman Sea (defined as east of 142°E). Note that the RMP results have not been updated for returns in 2005/2006 and beyond.

		Age at recapture						
		2	3	4	5	6	7	8
1990s RMP	All longline returns	39.4	46.7	53.5	58.6	60.3	64.1	45.7
	Japanese returns	39.1	48.1	50.9	54.2	56.5	59.3	42.9
2000s SRP	All longline returns	11.1	7.2	7.6	1.0	21.7	7.1	0.0
	Japanese returns	28.9	20.2	21.6	3.7	33.3	0.0	0.0

Table 7: Percent of (reported) longline recaptures from the Tasman Sea (defined as east of 142°E) that occurred in the Australian Fishing Zone (AFZ) for the 1990s RMP tag releases.

		Age at recapture						
		2	3	4	5	6	7	8
% in AFZ		91.2	77.7	68.2	51.6	42.8	35.1	21.6

Table 8: Estimates of age-specific fishing mortality rates (F) for different cohorts derived from tags released in the waters around southern Australia. Results are presented separately for tags released at different ages. Lower and upper 5% refer to the lower and upper 5th percentiles from the bootstrap estimates (i.e. the 90% confidence interval). Results were derived using reporting rate vector 1 from Table 1 for the surface fishery, group shedding rates from Table 2, natural mortality vector 1 from Table 3, and a reporting rate of 0.65 for the longline fisheries. (F estimates > 0.4 are shaded.)

Cohort	Age at release	Number of releases	Age	Year	F	Lower 5%	Upper 5%			
1999	2	750	2	2001	0.000	0.000	0.000			
			3	2002	0.036	0.020	0.057			
			4	2003	0.259	0.199	0.328			
			5	2004	0.099	0.050	0.159			
			6	2005	0.063	0.000	0.135			
			7	2006	0.041	0.000	0.097			
			8	2007	0.021	0.000	0.064			
			2000	1	1921	1	2001	0.000	0.000	0.000
2	2002	0.006				0.001	0.010			
3	2003	0.174				0.141	0.208			
4	2004	0.072				0.047	0.101			
5	2005	0.021				0.005	0.040			
6	2006	0.012				0.000	0.036			
7	2007	0.000				0.000	0.000			
2000	2	492				2	2002	0.004	0.000	0.011
			3	2003	0.272	0.207	0.348			
			4	2004	0.477	0.334	0.654			
			5	2005	0.453	0.227	0.826			
			6	2006	0.038	0.000	0.133			
			7	2007	0.175	0.000	0.582			
			2000	3	3276	3	2003	0.154	0.139	0.170
						4	2004	0.282	0.251	0.313
5	2005	0.156				0.124	0.190			
6	2006	0.079				0.049	0.112			
7	2007	0.017				0.004	0.033			
2001	1	2748				1	2002	0.000	0.000	0.000
						2	2003	0.008	0.004	0.012
						3	2004	0.215	0.181	0.252
			4	2005	0.057	0.036	0.078			
			5	2006	0.031	0.012	0.054			
			6	2007	0.015	0.004	0.033			
			2001	2	5869	2	2003	0.009	0.007	0.012
						3	2004	0.746	0.698	0.797
4	2005	0.685				0.597	0.806			
5	2006	0.550				0.410	0.754			
6	2007	0.051				0.018	0.096			
2001	3	1146				3	2004	0.583	0.512	0.662
						4	2005	0.789	0.632	1.009
						5	2006	1.491	0.822	3.063
			6	2007	0.591	-1.026	1.743			
			2002	1	3316	1	2003	0.001	0.000	0.002
						2	2004	0.023	0.016	0.030
						3	2005	0.115	0.092	0.139
						4	2006	0.075	0.050	0.103
5	2007	0.015				0.004	0.029			

2002	2	6256	2	2004	0.029	0.024	0.034
			3	2005	0.518	0.479	0.559
			4	2006	0.836	0.728	0.964
			5	2007	0.631	0.481	0.845
2002	3	720	3	2005	0.205	0.157	0.256
			4	2006	1.140	0.833	1.561
			5	2007	0.765	0.409	1.852
2003	1	2662	1	2004	0.000	0.000	0.000
			2	2005	0.044	0.031	0.057
			3	2006	0.502	0.425	0.587
			4	2007	0.363	0.281	0.467
2003	2	8692	2	2005	0.028	0.024	0.033
			3	2006	1.046	0.966	1.131
			4	2007	2.124	1.582	3.306
2003	3	3127	3	2006	0.285	0.252	0.321
			4	2007	0.491	0.433	0.557
2004	1	7084	1	2005	0.000	0.000	0.001
			2	2006	0.018	0.013	0.024
			3	2007	0.091	0.077	0.106
2004	2	7591	2	2006	0.025	0.020	0.030
			3	2007	0.262	0.242	0.282
2004	3	479	3	2007	0.168	0.123	0.224

Table 9: Comparison of age-specific fishing mortality rate estimates for a range of scenarios, as described below. (F estimates > 0.4 are shaded.)

Scenario 1: reporting rate vector 1 from Table 1 for the surface fishery, natural mortality vector 1 from Table 3, and a reporting rate of 0.65 for the longline fisheries (i.e., same as Table 8).

Scenario 2: same as scenario 1 but with reporting rate vector 4 from Table 1 for the surface fishery.

Scenario 3: same as scenario 1 but with a reporting rate of 0.30 for the longline fisheries.

Scenario 4: same as scenario 1 but with natural mortality vector 2 from Table 3.

Cohort	Age at release	Number of releases			F estimate						
			Age	Year	Scenario 1	Scenario 2	Scenario 3	Scenario 4			
1999	2	750	2	2001	0.000	0.000	0.000	0.000			
			3	2002	0.036	0.044	0.058	0.039			
			4	2003	0.259	0.372	0.325	0.301			
			5	2004	0.099	0.120	0.177	0.127			
			6	2005	0.063	0.064	0.074	0.084			
			7	2006	0.041	0.047	0.108	0.058			
			8	2007	0.021	0.025	0.060	0.031			
			2000	1	1921	1	2001	0.000	0.000	0.000	0.000
2	2002	0.006				0.007	0.009	0.006			
3	2003	0.174				0.249	0.198	0.205			
4	2004	0.072				0.085	0.100	0.093			
5	2005	0.021				0.023	0.050	0.030			
6	2006	0.012				0.009	0.013	0.017			
7	2007	0.000				0.000	0.000	0.000			
2000	2	492				2	2002	0.004	0.004	0.008	0.004
			3	2003	0.272	0.397	0.315	0.295			
			4	2004	0.477	0.644	0.591	0.582			
			5	2005	0.453	0.584	0.776	0.682			
			6	2006	0.038	0.059	0.142	0.070			
			7	2007	0.175	0.279	0.606	0.362			
			2000	3	3276	3	2003	0.154	0.224	0.163	0.154
						4	2004	0.282	0.345	0.313	0.302
5	2005	0.156				0.159	0.178	0.179			
6	2006	0.079				0.067	0.107	0.096			
7	2007	0.017				0.019	0.033	0.022			
2001	1	2748				1	2002	0.000	0.000	0.000	0.000
						2	2003	0.008	0.010	0.011	0.009
						3	2004	0.215	0.242	0.236	0.255
			4	2005	0.057	0.053	0.072	0.074			
			5	2006	0.031	0.026	0.050	0.043			
			6	2007	0.015	0.015	0.034	0.022			
			2001	2	5869	2	2003	0.009	0.011	0.015	0.009
						3	2004	0.746	0.879	0.833	0.829
4	2005	0.685				0.704	0.919	0.947			
5	2006	0.550				0.478	1.547	1.254			
6	2007	0.051				0.054	0.326	0.194			
2001	3	1146				3	2004	0.583	0.674	0.648	0.585
						4	2005	0.789	0.752	0.930	0.866
						5	2006	1.491	0.877	NA	2.715
			6	2007	0.591	0.283	-1.092	NA			
			2002	1	3316	1	2003	0.001	0.001	0.001	0.001
						2	2004	0.023	0.025	0.029	0.025
						3	2005	0.115	0.102	0.128	0.135
						4	2006	0.075	0.054	0.094	0.096

			5	2007	0.015	0.014	0.016	0.021
2002	2	6256	2	2004	0.029	0.032	0.034	0.029
			3	2005	0.518	0.445	0.561	0.568
			4	2006	0.836	0.477	1.165	1.131
			5	2007	0.631	0.342	1.492	1.545
2002	3	720	3	2005	0.205	0.180	0.227	0.206
			4	2006	1.140	0.606	1.290	1.279
			5	2007	0.765	0.342	1.078	1.150
2003	1	2662	1	2004	0.000	0.000	0.000	0.000
			2	2005	0.044	0.040	0.053	0.048
			3	2006	0.502	0.321	0.571	0.617
			4	2007	0.363	0.273	0.416	0.547
2003	2	8692	2	2005	0.028	0.025	0.035	0.028
			3	2006	1.046	0.594	1.193	1.189
			4	2007	2.124	0.751	NA	NA
2003	3	3127	3	2006	0.285	0.187	0.303	0.285
			4	2007	0.491	0.404	0.523	0.530
2004	1	7084	1	2005	0.000	0.000	0.001	0.000
			2	2006	0.018	0.014	0.023	0.020
			3	2007	0.091	0.086	0.106	0.107
2004	2	7591	2	2006	0.025	0.019	0.033	0.025
			3	2007	0.262	0.245	0.285	0.284
2004	3	479	3	2007	0.168	0.159	0.189	0.169

Table 10: First year recaptures of fish released in December in the Great Australian Bight by age of release and fishing year. Estimated percent caught is based on weighted reporting rate estimates in Table 1, with no allowance for tag shedding.

Fishing year	Release age	Number released	Number returned	Percent returned	Est. percent caught
2003	1	17	0	0	0
	2	894	19	2.1	3.3
	3	3004	295	9.8	15.3
	4	242	34	14.0	22.0
	5	8	3	37.5	58.6
2004	1	622	0	0	0
	2	3187	82	2.6	5.1
	3	978	251	25.7	51.0
	4	27	7	25.9	51.5
	5	3	0	0	0
2005	1	52	0	0	0
	2	2760	43	1.6	3.9
	3	308	34	11.0	27.9
	4	130	12	9.2	23.3
	5	68	12	17.6	44.6
2006	1	22	0	0	0
	2	1887	20	1.1	3.5
	3	2442	193	7.9	26.1
	4	14	1	7.1	23.6
	5	1	0	0	0
2007	1	5	0	0	0
	2	3023	51	1.7	4.0
	3	281	24	8.5	20.1
	4	0	0	-	-
	5	0	0	-	-

Figure 1: Comparison of fishing mortality rate (F) estimates at ages 3 and 4 obtained from the SRP tagging with those obtained from the 1990s RMP tagging. Estimates for the 1990s tagging were obtained using a Brownie model that integrates data from all release ages.

