



**Australian Government**  
**Department of Agriculture**  
ABARES

# **Research mortality allowance**

## **Proposed allowance for 2015 and 2014**

### **usage report**

S. Hansen and H. Patterson

Research by the Australian Bureau of Agricultural  
and Resource Economics and Sciences

August 2014



Working Paper CCSBT-ESC/1409/28 prepared for the CCSBT  
Extended Scientific Committee for the 19<sup>th</sup> Meeting to the Scientific  
Committee

1–6 September 2014, Auckland, New Zealand

### **Ownership of intellectual property rights**

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

### **Creative Commons licence**

All material in this publication is licensed under a Creative Commons Attribution 3.0 Australia Licence, save for content supplied by third parties, logos and the Commonwealth Coat of Arms.



Creative Commons Attribution 3.0 Australia Licence is a standard form licence agreement that allows you to copy, distribute, transmit and adapt this publication provided you attribute the work. A summary of the licence terms is available from [creativecommons.org/licenses/by/3.0/au/deed.en](http://creativecommons.org/licenses/by/3.0/au/deed.en). The full licence terms are available from [creativecommons.org/licenses/by/3.0/au/legalcode](http://creativecommons.org/licenses/by/3.0/au/legalcode).

### **Cataloguing data**

Hansen, S & Patterson, H 2014, Research Mortality Allowance – Proposed allowance for 2015 and 2014 usage report, ABARES, Canberra, August. CC BY 3.0.

### **Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES)**

Postal address GPO Box 858 Canberra ACT 2601

Switchboard +61 2 6272 2010

Facsimile +61 2 6272 2001

Email [info.abares@daff.gov.au](mailto:info.abares@daff.gov.au)

Web [daff.gov.au/abares](http://daff.gov.au/abares)

Inquiries about the licence and any use of this document should be sent to [copyright@daff.gov.au](mailto:copyright@daff.gov.au).

The Australian Government acting through the Department of Agriculture, represented by the Australian Bureau of Agricultural and Resource Economics and Sciences, has exercised due care and skill in preparing and compiling the information and data in this publication. Notwithstanding, the Department of Agriculture, ABARES, its employees and advisers disclaim all liability, including for negligence and for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying upon information or data in this publication to the maximum extent permitted by law.

### **Acknowledgements**

Thank you to CSIRO, SARDI and University of Tasmania for providing information on the projects and to SBTMAC for comments on proposals. Ilona Stobutzki provided comments on the paper.

# Contents

Summary	1
<b>1 Proposals for 2015</b>	<b>2</b>
i) Electronic tagging and effect of seismic exploration	2
ii) Does the "membrane pacemaker" theory hold true for endothermic fish?	3
iii) Status, distribution and abundance of iconic species and apex predators in the Great Australian Bight	4
iv) Health assessment of wild southern bluefin tuna	4
<b>2 Summary of 2014 RMA usage</b>	<b>6</b>
References	7

# Summary

Australia requests research mortality allowances (RMA) for four projects on southern bluefin tuna (SBT) in 2015. Most of the projects aim to avoid SBT mortality and are requesting RMA to cover any incidental mortality. The RMA amounts requested are relatively small and appropriate for the individual projects. There are four projects requesting RMA that are described in this paper:

- 1) A 3 t RMA is requested to continue initiatives focused on investigating: (a) the spatial dynamics and mortality rates of SBT utilising electronic tagging techniques; (b) the impacts of noise associated with oil and gas exploration on SBT in the Great Australian Bight. Note that RMA has been granted previously for this project and is to cover incidental mortalities only. No RMA for this project has been used in recent years.
- 2) A 0.5 t RMA is requested for a study examining the molecular basis for endothermy using SBT as a novel model. This project will involve the capture of small SBT off southern Western Australia. Larger fish will be purchased from Port Lincoln aquaculture farms.
- 3) A 1.25 t RMA is requested for a project examining iconic and apex predator species in the Great Australian Bight as part of a larger ecosystem project. This project will involve some research fishing activities that may incidentally capture SBT. Note that RMA has been granted previously for this project and is to cover incidental mortalities only. No RMA for this project has been used in 2014 (to date).
- 4) Finally, a 1.2 t RMA is requested for a project examining the health of wild SBT. Note that RMA has been granted previously for this project. A total of 663 kg of SBT was used for this project in 2014.

The overall RMA requested by Australia is small (5.95 t) and is for projects that will enhance our understanding of the biology and ecology of SBT.

In 2014, Australia was granted 5.95 t of RMA for four projects (three of those projects are requesting RMA for 2015). To date, two projects have used RMA, totalling 1.1 t.

# 1 Proposals for 2015

## i) Electronic tagging and effect of seismic exploration

The major aim of this project is to continue previous tagging initiatives aimed at examining spatial dynamics (movements, mixing, residency, regional fidelity and spawning frequency) of wild SBT. This study involves the tagging of SBT across a range of sizes (25–100 kg) with archival and pop-up satellite archival tags throughout the south-eastern Australian region in 2015, as well as conducting controlled exposure experiments investigating the impacts of seismic exploration on SBT behaviour and anatomy.

Tagging efforts to date have been critical in improving our understanding of the spatial dynamics and habitat use of SBT and are fundamental to the ongoing spatial management of the domestic fishery. Use of similar, and newer technology (mini-PATs) to that used previously, in this next phase of research will allow us to add to current understanding of the spatial dynamics of juvenile and adult SBT and better understand the spatial dynamics of medium size ranges of SBT (30+ kg), data which are currently lacking. Continued tagging of larger size classes will allow for better understanding and characterization of spawning migrations, data which are relevant to the parameterisation of spawning stock biomass (SSB) in the operating model. Success of tagging initiatives to date suggests that releases of electronic tags on SBT are feasible and cost effective.

The aim of this project, undertaken by scientists at the Commonwealth Scientific and Industrial Research Organisation (CSIRO), is to continue tagging initiatives. This study is proposing to release in the order of 150–200 tags on SBT ranging in size from 20 kg to 100+ kg. The majority of fish are planned to be tagged in 2014–15, with any tags not deployed in 2014–15, deployed the following year 2015–16. Approval for 2014 was applied for under CCSBT-ESC/1309/21 and approved (see agenda item 14(153) of Anon (2013)).

The study will also investigate the impacts of noise associated with oil and gas exploration on SBT in the Great Australian Bight (GAB). The GAB is particularly important for young SBT (1–4 years), with large numbers migrating into the warm, shelf waters each summer to feed on abundant prey species. Scientific aerial surveys for juvenile SBT in the GAB collect data on surface schools of SBT observed from light planes that fly designated line transects according to established and consistent protocols. These data are used to estimate an annual index of relative abundance of 2–4 year olds in the GAB between December and March (Eveson et al. 2012). This index forms one of two relative abundance series used directly as an input data series for the Management Procedure used by the CCSBT to recommend the global TAC (Anon 2013).

There is concern that activities associated with oil and gas exploration, specifically seismic surveys and noise associated with exploratory drilling activities, may impact on the migration and behaviour of SBT in the GAB. A pilot controlled exposure experiment developed in consultation with industry in Port Lincoln, and done in collaboration with Curtin University and research staff of the Australian Southern Bluefin Tuna Industry Association is proposed. A small number of juvenile SBT in experimental cages will be exposed to a range of levels of seismic noise representative of that used in oil and gas exploration in the GAB and compared with SBT in "control" cages (i.e. not exposed to direct seismic noise). A range of technologies including stomach temperature sensors and acoustic tags will be used to monitor the feeding and swimming behaviour of individuals. Weights and lengths of individuals, along with amount of feed provided will be monitored, to determine if growth rates and feed conversion ratios at the

end of the trial deviated from expected rates of growth. Necropsies of caged fish will be conducted to assess any immediate post-exposure organ damage and also later in the trial to examine rates of recovery.

This proposal extends the 3 t RMA request granted for 2014, to cover SBT tagging in 2015, noting that none of the RMA requested for 2014 has been used as yet (as of July 2014), due to delays in funding arrangements.

Results of earlier initiatives investigating the spatial dynamics of SBT are provided in Gunn and Patterson (2003), Patterson et al. (2005), Gunn et al. (2006), Evans and Patterson (2007), Patterson et al. (2008), Evans et al. (2012) and Basson et al. (2012). Data from these projects are integral in spatial management procedures of the Australian Fishery implemented by the Australian Fisheries Management Authority. Use of these data in spatial management is detailed in Hobday and Hartmann (2006), Hobday et al. (2011) and Hartog et al. (2011).

## **ii) Does the "membrane pacemaker" theory hold true for endothermic fish?**

This project, undertaken by CSIRO, will use SBT as a model to test prevailing theories of the evolution of endothermy (warm-bloodedness). Bluefin tunas are top predators whose removal or displacement due to over-fishing or climate change could destabilize marine ecosystems. Captive breeding is a possible solution but its success is limited by our inadequate understanding of tuna thermal biology. This project will rectify this.

This project will investigate the molecular basis of endothermy using a novel model, SBT. Endotherms maintain their internal body temperature by conserving metabolically generated heat, whereas ectotherms depend on heat from the environment. Despite its evolutionary significance, the molecular basis of endothermy is still not well understood. Tunas provide unique models to investigate this because their bodies consist of both endothermic and ectothermic organs and they only develop the capacity for endothermy during the first year of their lives. The outcome of this project will be a greater understanding of vertebrate thermal biology leading to better predictions of the effects of climate change on ecosystem stability.

Recently it was suggested that the limited success of bluefin tuna captive breeding may be due to under-developed capacity for endothermy when the young fish are transferred from the warm water of the hatchery to the cooler water of the sea cages. This was based on observations of Pacific bluefin tuna but it may also be true for SBT. The growth of the bluefin tuna aquaculture industry is limited by high rates of cold-induced mortalities in captive bred fish proposed to be due to underdeveloped endothermy. Small (incipiently endothermic) southern bluefin tuna are found in Australian coastal waters from Western Australia to the GAB.

We will compare small (40 to 60 cm fork length) SBT with similarly sized Australian bonito, a close ectothermic relative. These species, in this size range, are found in waters off southern Western Australia during the period November to February each year. We seek to capture small SBT in waters between Albany and Fremantle. Larger fish will be purchased from the Port Lincoln aquaculture businesses.

The specific aims of this project are:

- 1) To investigate the ontogeny of red muscle endothermy in small (incipiently endothermic) SBT.

- 2) To compare mitochondrial respiration rate, proton leak rate and membrane lipid unsaturation in endothermic versus ectothermic organs/tissues of large (fully endothermic) SBT and similarly-sized Australian bonito.
- 3) To determine the effects of varying membrane unsaturation on mitochondrial respiration rate and proton leak rate in a recently established SBT cell line.
- 4) To explore the transcriptional regulation of mitochondrial biogenesis and its role in the ontogeny of red muscle endothermy in fish.

An RMA of 0.5t is requested for this study, representing lethal sampling of juvenile SBT.

### **iii) Status, distribution and abundance of iconic species and apex predators in the Great Australian Bight**

The project, undertaken by researchers at the South Australian Research and Development Institute, is part of a larger project investigating aspects of the ecosystem of the GAB. The aims of this portion of the project are to provide background information on the relative distributions and abundances of key iconic and apex predator species in the GAB. It has four main tasks including:

- 1) Assessing the occurrence and distribution of pygmy blue whales and other cetaceans using offshore habitats of the GAB;
- 2) Assessing the occurrence and distribution of southern right whales and other cetaceans using inshore habitats of the GAB;
- 3) Obtaining abundance indices of key pinniped and seabird populations in the GAB; and
- 4) Characterising the spatial and temporal distribution of pelagic sharks and large teleosts in the GAB.

We estimate the capture of a maximum of 50 SBT from pelagic longline sets in the east GAB region off South Australia. Task 4 of this study involves sampling with pelagic longlines (for electronic tagging) in the eastern GAB and may therefore catch SBT and result in some mortalities. Therefore, the proposal requests 1.25 t of RMA for 2015.

This proposal extends the 1.25 t RMA request granted for 2014. However, the project commencement was delayed, and no RMA has been used in 2014 (as of July 2014).

### **iv) Health assessment of wild southern bluefin tuna**

The major aim of this project is to assess the health of wild SBT. As well as traditional health assessments including histology, microbiology, immunology and haematology, this project will aim to further develop and apply new molecular methods, including those for pathogen detection and associated pathology in wild SBT. New molecular methods will be developed which could help to identify specific blood borne biomarkers for various pathogens (disease agents) and ultimately could lead to non lethal health assessment for these animals.

This project, undertaken by researchers from the University of Tasmania, will have significant scientific and ethical benefits. The results of this project will greatly contribute to the current lack of knowledge of the health of wild SBT populations including pathogen prevalence and/or intensities. Identification of suitable health biomarkers could also negate the need for destructive sampling of animals. This project could therefore have implications for future SBT research.

Most health assessments for SBT have been conducted using ranched animals. Comparatively, the health status of non-ranched wild SBT has received little attention. Some wild SBT sampling has previously been conducted with participation of SBT industry as part of Aquafin Cooperative Research Centre and Fisheries Research and Development Corporation projects. This research was based on frozen and formalin fixed samples that were collected by the industry from tuna schools captured for ranching so for example the samples for pathogen detection (particularly some species of parasites) were limited. All SBT were from GAB, further limiting interpretation of the results. Some of the results from this work with wild tuna have been published (please see Rough et al. 2005, Aiken et al. 2007, Aiken et al. 2008, Aiken et al. 2009) and one manuscript has recently been accepted for publication (Kirchhoff et al. 2014).

This proposal extends the 1.2 t RMA request granted for 2014, to cover further work in 2015. This request is based on an estimate of 20 SBT with a maximum weight of 60 kg each. In 2014, a total of 30 SBT were caught for this project, amounting to an estimated 663 kg.

To date, publications from this research have not yet been completed, the first papers are anticipated to be submitted for publication later this year. The papers will be provided to the ESC for information once available.



## 2 Summary of 2014 RMA usage

A total of 5.95 t of RMA was granted in 2013 for Australian research projects undertaken in 2014. To date (as of July 2014), 1.1 t of RMA has been utilised in 2014, well below the amount granted, although the possibility remains that tagging will commence in the first project listed below in December 2014.

- 1) A 3 t RMA was approved to continue initiatives focused on investigating: (a) the spatial dynamics and mortality rates of SBT utilising electronic tagging techniques; (b) the impacts of noise associated with oil and gas exploration on SBT in the Great Australian Bight. Note that RMA is to cover incidental mortalities only. No RMA for this project has been used in recent years.
- 2) A 1.25 t RMA was approved for a project examining iconic and apex predator species in the Great Australian Bight as part of a larger ecosystem project. To date, there has been no SBT mortality associated with this project in 2014.
- 3) An RMA of 1.2 t was approved for a project examining the health of wild SBT. A total of 663 kg of SBT was used for this project in 2014. To date, there are no publications generated from this research, the first papers are anticipated to be submitted for publication later this year. These publications will be provided to the ESC once available.
- 4) Finally, 0.5 t of RMA was approved for a study examining the post-release survival of SBT from recreational fishing. This RMA was requested for incidental mortality only. There was 451 kg of SBT mortality from this study in 2014. Incidental mortality was largely associated with seal strike (153 kg landed after seal attack, and approximately 260 kg lost during capture process [based on a 20 kg per fish average]). This research was conducted as part of a Fisheries Research and Development Corporation project (No. 2013/025). The report for this work is expected to be available mid-2015 and will be provided to the ESC.

# References

- Aiken, H, Hayward, CJ, Cameron, A & Nowak, BF 2009, 'Simulating blood fluke, *Cardicola forsteri*, infection in farmed southern bluefin tuna, *Thunnus maccoyi*, using stochastic models. *Aquaculture*, vol. 293, pp. 204-210.
- Aiken, HM, Bott, NJ, Mladineo, I, Montero, FE, Nowak, BF & Hayward, CJ 2007, 'Molecular evidence for cosmopolitan distribution of platyhelminth parasites of tunas (*Thunnus* spp.)' *Fish and Fisheries Science*, vol. 8, pp. 167-180.
- Aiken, HM, Hayward, CJ, Crosbie, PB, Watts, M & Nowak, BF 2008, 'Serological evidence of an antibody response in farmed southern bluefin tuna naturally infected with the blood fluke *Cardicola forsteri*' *Fish and Shellfish Immunology*, vol. 25, pp. 66-75.
- Anon 2013, 'Report of the eighteenth meeting of the CCSBT Scientific Committee' Canberra, Australia, 7 September 2013.
- Basson, M, Hobday, AJ, Eveson, JP & Patterson, TA 2012, 'Spatial interactions among juvenile southern bluefin tuna at the global scale: A large scale archival tag experiment, FRDC Report 2003/002.
- Evans, K & Patterson, T 2007, 'Movements and behaviour of large SBT in the Australian region determined using pop-up archival satellite tags: a summary of results for 2006-07' CCSBT-ESC/0709/Info 01, Hobart, Australia, 10-14 September 2007.
- Evans K, Patterson TA, Reid H and Harley SJ 2012, 'Reproductive schedules in southern bluefin tuna: are current assumptions appropriate?' *PLoS ONE*, 7: e34550.
- Eveson P, Farley J and Bravington M 2012, 'The aerial survey index of abundance: updated analysis methods and results for the 2011/12 fishing season' CCSBT-ESC/1208/16, Tokyo, Japan, 27-31 August 2012.
- Gunn J and Patterson T 2003, 'A pilot study to examine the potential for using pop-up satellite transmitting archival tags (PATs) to examine the migrations and behavior of adult Southern Bluefin Tuna (SBT)' CCSBT-ESC/0309/Info 2, Christchurch, New Zealand, 1-4 September 2003.
- Gunn J, Evans K, Patterson T and Carter T 2006, 'Examining the movement and residency of adult SBT in the Tasman Sea and on their spawning grounds south of Indonesia using pop-up archival tags' CCSBT-ESC/0609/Info 01, Miyazaki, Japan, 10-13 October 2006.
- Hartog, J, Hobday, AJ, Matear, R & Feng, M 2011, 'Habitat overlap of southern bluefin tuna and yellowfin tuna in the east coast longline fishery - implications for present and future spatial management' *Deep Sea Research Part II*, vol 58, pp. 746-752.
- Hobday, AJ & Hartmann, K 2006, Near real-time spatial management based on habitat predictions for a longline bycatch species. *Fisheries Management and Ecology*, vol. 13, pp. 365-380.
- Hobday, AJ, Hartog, JR, Timmis, T & Fielding, J 2010, 'Dynamic spatial zoning to manage southern bluefin tuna capture in a multi-species longline fishery' *Fisheries Oceanography*, vol. 19, pp. 243-253.

Kirchhoff, NT, Nelligan, J, Ellis, D, Nosworthy, E, Cadoret, K, Leef, M, & Nowak, B 2014, 'Inter-annual and intra-annual variability in blood variables and parasitic loads of wild *Thunnus maccoyii*' *Canadian Journal of Fisheries and Aquatic Science*, in press.

Patterson, TA, Evans, K, Carter, TI & Gunn, JS 2008, 'Movement and behaviour of large southern bluefin tuna (*Thunnus maccoyii*) in the Australian region determined using pop-up satellite archival tags' *Fisheries Oceanography*, vol. 17, pp. 352–367.

Patterson, T, Gunn, J, Evans, K & Carter, T 2005, 'Movement and residency of adult SBT in the Tasman Sea and on their spawning grounds south of Indonesia using pop-up archival tags: a summary of results for 2004' CCSBT-ESC/0409/29, Jeju, Korea, 13–16 September 2006.

Rough, KM, Nowak, BF & Reuter, R 2005, 'Haematology and leucocyte morphology of wild caught SBT' *Journal of Fish Biology*, vol. 66, pp. 1649-1659.