



## INTRODUCTION

There are many species of seabirds that have been recorded as bycatch in pelagic longline fisheries. The majority of these, both in terms of species and numbers caught, are albatrosses and petrels (Anderson *et al.* 2011; Ryan *et al.* 2002). Of the 22 species of albatrosses found globally, 18 overlap in distribution with the Southern Bluefin Tuna (SBT) fishery, as do seven of the eight petrel species also listed under the Agreement on the Conservation of Albatrosses and Petrels (ACAP). This paper provides a summary of the status and trends of albatross and petrel species listed under Annex 1 of ACAP that breed or forage in areas where Southern Bluefin tuna are fished.

## CONSERVATION STATUS

Of the 18 species of albatrosses with distributions that overlap with the SBT fishery, the International Union for Conservation of Nature (IUCN) lists:

- two as *Critically Endangered* (CR),
- five as *Endangered* (EN),
- six as *Vulnerable* (VU) and
- five as *Near Threatened* (NT).

Of the seven ACAP petrel species with distributions that overlap with the SBT fishery, the International Union for Conservation of Nature (IUCN) lists:

- four are listed as VU,
- one as NT and
- two as *Least Concern* (LC).

The IUCN status of two albatross species has changed since ERSWG 10 in 2013. The Black-browed albatross *Thalassarche melanophris* has been downlisted from EN to NT, and the Grey-headed albatross *Thalassarche chrysostoma* has been uplisted from VU to EN. Both of these changes were based on improved knowledge, rather than genuine improvement or deterioration in status.

## STATUS OF KNOWLEDGE RELATING TO POPULATION SIZE AND TRENDS

Comprehensive knowledge of population size, trend and demographic parameters are fundamental to many aspects of albatross and petrel conservation, and vital to monitoring the effectiveness of management actions. Although the size of most populations has been determined at some point in time, the trend and current demographic statistics for many populations are not known, due to the high level of resources required to access remote sites at appropriate intervals. Determination of global trends can also be difficult because populations of the same species at different sites may show different trajectories.

Estimates of bycatch in global longline fisheries indicate that 160,000 – 320,000 seabirds, mostly albatrosses, petrels and shearwaters, are killed each year (Anderson *et al.* 2011). These estimates may be understated by as much as 50% or more due to lack of observer data, under-estimates in observer data, or under-reporting (Brothers *et al.* 2010, Anderson *et al.* 2011). Such levels of incidental mortality are known to have resulted in or are contributing to population declines for a number of these species (e.g. Wanless *et al.* 2009; Weimerskirch *et al.* 1997).

At its second meeting in September 2014, ACAP's Population and Conservation Status Working Group examined the current (1993-2013) global trends of species listed under the Agreement (**Table 1**). The approach combines census information submitted to the ACAP database ([data.acap.aq](http://data.acap.aq)) and results of published population models. The time span of two decades was considered appropriate to reflect the trend of these long lived species, some of which breed only every two years, and which may show high annual variation in breeding numbers. The confidence of the assigned trend reflects both the accuracy and extent of the population data. The Working Group envisages that the trends will be reviewed on a triennial basis or sooner if significant new information becomes available for any of the species.

Further information can also be found in the species assessments developed by ACAP ([www.acap.aq/index.php/species-assessments](http://www.acap.aq/index.php/species-assessments)) which provide comprehensive information on the distribution, biology and threats facing all ACAP species.

## REFERENCES

- Anderson, O. R. J., Small, C. J., Croxall, J. P., Dunn, E. K., Sullivan, B. J. Yates, O. and Black, A. 2011. Global seabird bycatch in longline fisheries. *Endangered Species Research* **14**: 91-106.
- Brothers, N., Duckworth, A.R., Safina, C. and Gilman, E.L. 2010. Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PLoS ONE* **5** (8): e12491. Doi10.1371/journal.pone.0012491.
- Ryan, R.G., Keith, D.G. and Kroese, M. 2002. Seabird bycatch by tuna longline fisheries off southern Africa, 1998-2000. *South African Journal of Marine Science* **24**: 103-110.
- Wanless, R. M., Ryan, P.G, Altwegg, R., Angel, A., Cooper, J., Cuthbert, R., and Hilton, G.M. 2009. From both sides: Dire demographic consequences of carnivorous mice and longlining for the Critically Endangered Tristan albatrosses on Gough Island. *Biological Conservation* **142**: 1710-1718.

Weimerskirch, H. N., Brothers, N., and Jouventin, P. 1997. Population dynamics of wandering albatross *Diomedea exulans* and Amsterdam albatross *D. amsterdamensis* in the Indian Ocean and their relationships with long-line fisheries: Conservation implications. *Biological Conservation* **79**: 257-270.

Table 1. 2014 summary of status and trends of ACAP species breeding/foraging in the SBT fishery area.

Common name	Scientific name	Number of sites <sup>1</sup>	Single Country Endemic	Breeding Frequency <sup>2</sup>	Annual breeding pairs (latest census year) <sup>1</sup>	Current Trend 1993-2013 <sup>3</sup>	Trend Confidence	IUCN Status 2014 <sup>4</sup>	IUCN Status Justification <sup>4</sup>
Amsterdam Albatross	<i>Diomedea amsterdamensis</i>	1	France	B	31 (2013)	↑	High	CR	Extremely small population, confined to a tiny area on one island. Although numbers have recently been increasing, a continuing decline is projected owing to the impact of a disease which is probably already causing chick mortality.
Tristan Albatross	<i>Diomedea dabbenena</i>	1	UK	B	1,650 (2014)	↓	High	CR	Extremely small breeding range and a projected extremely rapid population decline over three generations (70 years). Modelled population declines are a consequence of very low adult survival owing to incidental mortality in longline fisheries, compounded by low fledging success caused by predation of chicks by introduced mice.
Atlantic yellow-nosed Albatross	<i>Thalassarche chlororhynchos</i>	6	UK	A	33,650 (1974-2011)	↔	Low	EN	Very small breeding range and is estimated to be undergoing a very rapid ongoing decline projected over three generations (72 years) owing to incidental mortality in longline fisheries.
Grey-headed Albatross	<i>Thalassarche chrysostoma</i>	29		B	97,716 (1982-2014)	↓	Medium	EN	Uplisted to Endangered as data from some major colonies suggest that overall declines are taking place at a very rapid rate over three generations (90 years), even if colonies lacking trend information are assumed to be stable. The major driver of declines is likely to be incidental mortality on longline fisheries.
Indian yellow-nosed Albatross	<i>Thalassarche carteri</i>	6		A	39,320 (1984-2009)	↓	Medium	EN	Estimated very rapid ongoing decline over three generations (71 years), based on data from the population stronghold on Amsterdam Island. This decline is the result of adult mortality and poor recruitment owing to interactions with fisheries and disease.
Northern royal Albatross	<i>Diomedea sanfordi</i>	5	NZ	B	5,782 (1995-2013)	?	-	EN	Restricted to a tiny breeding range in which severe storms in the 1980s resulted in a decrease in habitat quality, which led to poor breeding success. Based on this low breeding success, the population is estimated and projected to be undergoing a very rapid decline over three generations (1985-2069). Evidence suggests that the number of breeding pairs may have remained relatively stable, thus the species might qualify for downlisting in the future, however in the absence of recent substantive data upon which to assess trends or changes in productivity rates, projected declines are precautionarily maintained.

Common name	Scientific name	Number of sites <sup>1</sup>	Single Country Endemic	Breeding Frequency <sup>2</sup>	Annual breeding pairs (latest census year) <sup>1</sup>	Current Trend 1993-2013 <sup>3</sup>	Trend Confidence	IUCN Status 2014 <sup>4</sup>	IUCN Status Justification <sup>4</sup>
Sooty Albatross	<i>Phoebastria fusca</i>	15		B	12,170 (1974-2014)	↓	Very Low	EN	Very rapid decline over three generations (90 years), probably owing to interactions with fisheries. Since 1980, three sites (Crozet, Marion and Gough) have witnessed severe declines, although the population at Prince Edward may have increased between 2002-2009. However, high variability in population counts between years necessitates caution and further data are required before a change in status should be considered.
Antipodean Albatross	<i>Diomedea antipodensis</i>	6	NZ	B	8,274 (1995-2010)	↓	Medium	VU	Largely confined to three small islands when breeding and is therefore highly susceptible to stochastic effects and human impacts. Recent data (2005-2008) from the Auckland Islands indicate declines in adult survival, productivity and recruitment, which, if confirmed by further monitoring, could result in a reclassification of EN or CR.
Black Petrel	<i>Procellaria parkinsoni</i>	2	NZ	A	1,577 (1998-2013)	↓	Medium	VU	Breeds on just two very small islands where introduced predators are a potential threat. The population is assumed to be stable, but if a decline is detected, the species should be uplisted to EN.
Campbell Albatross	<i>Thalassarche impavida</i>	2	NZ	A	21,648 (2012)	↔	Low	VU	Breeding is restricted to a single location, where it is susceptible to potential human impacts and stochastic events. Although numbers decreased steeply between the 1970s and 1980s owing to interactions with fisheries, the population is now thought to be increasing, although there has not been a census since 1996.
Chatham Albatross	<i>Thalassarche eremita</i>	1	NZ	A	5,245 (2011)	↔	Medium	VU	Very small breeding range (restricted to one breeding site) rendering it susceptible to stochastic events and human impacts.
Salvin's Albatross	<i>Thalassarche salvini</i>	12	NZ	A	42,219 (1986-2013)	↓	Low	VU	May have undergone a rapid decline, but different census methods make a comparison of the available data potentially misleading. However, breeding is largely restricted to one tiny island group, where it is susceptible to stochastic events.
Southern royal Albatross	<i>Diomedea epomophora</i>	4	NZ	B	7,941 (1991-2008)	↔	Medium	VU	Although current population trends are assumed to be stable, this species has a very small range, breeding on four islands, although largely confined to just one, therefore highly susceptible to stochastic effects and human impacts.
Spectacled Petrel	<i>Procellaria conspicillata</i>	1	UK	A	14,400 (2010)	↑	High	VU	Owing to its very small breeding range, it is highly susceptible to stochastic events and human activities. Any evidence of population declines would likely lead to its uplisting.

Common name	Scientific name	Number of sites <sup>1</sup>	Single Country Endemic	Breeding Frequency <sup>2</sup>	Annual breeding pairs (latest census year) <sup>1</sup>	Current Trend 1993-2013 <sup>3</sup>	Trend Confidence	IUCN Status 2014 <sup>4</sup>	IUCN Status Justification <sup>4</sup>
Wandering Albatross	<i>Diomedea exulans</i>	28		B	8,132 (1981-2013)	↓	High	VU	Overall, past and predicted future declines amount to a rapid population reduction over a period of three generations. This species is undergoing a rapid decline in the South Atlantic, as well as on the Crozet and Kerguelen Islands. Longline fishing is believed to be a main cause of decline, causing reductions in adult survival and juvenile recruitment, and this threat is ongoing.
Westland Petrel	<i>Procellaria westlandica</i>	1	NZ	A	2,827 (2011)	↔	Low	VU	Restricted to one very small area when breeding, rendering the population highly vulnerable to stochastic events and other potential threats.
White-chinned Petrel	<i>Procellaria aequinoctialis</i>	73		A	1,057,930 (1984-2012)	↓	Very Low	VU	Suspected rapid declines, although almost no reliable estimates of historical populations exist. Very high rates of incidental mortality in longline fisheries have been recorded in recent years; the probability that these circumstances will continue and its susceptibility to predation and the degradation of breeding habitat indicate that a rapid and on-going population decline is likely.
Black-browed Albatross	<i>Thalassarche melanophris</i>	65		A	673,048 (1982-2013)	↑	High	NT	Downlisted to Near Threatened as it is no longer estimated to be undergoing very rapid population declines. However there remains a considerable degree of uncertainty over population trends for a significant part of the global population, and trend estimates are heavily influenced by the extrapolation over 65 years of data from a ten-year period. In addition, high levels of mortality of this species are reported from longline and trawl fisheries in the South Atlantic. For these reasons, moderately rapid ongoing declines over three generations since 1980 are precautionarily suspected until further data are forthcoming.
Buller's Albatross	<i>Thalassarche bulleri</i>	10	NZ	A	29,948 (1971-2012)	↔	Low	NT	Although the species is restricted to a tiny small area when breeding, the population is stable and the islands on which it breeds are moderately widely spread.
Grey Petrel	<i>Procellaria cinerea</i>	17		A	79,649 (1981-2012)	↓	Very Low	NT	Although there are no current trend data, this species is susceptible to introduced mammalian predators, and today it is the most commonly caught bycatch species in longline fisheries in New Zealand waters. Evidence from Gough Island, formerly thought to contain the largest population of this species, suggest that the species is likely to be subjected to considerable predation from introduced mice that are a major predator on other winter-breeding seabirds. The population on the Kerguelen Islands may also be in decline due to fishery bycatch. Based on these data a moderately rapid decline is suspected, but further data are urgently required in order to more accurately assess its population numbers and trends.

Common name	Scientific name	Number of sites <sup>1</sup>	Single Country Endemic	Breeding Frequency <sup>2</sup>	Annual breeding pairs (latest census year) <sup>1</sup>	Current Trend 1993-2013 <sup>3</sup>	Trend Confidence	IUCN Status 2014 <sup>4</sup>	IUCN Status Justification <sup>4</sup>
Light-mantled Albatross	<i>Phoebastria palpebrata</i>	71		B	12,082 (1954-2013)	↔	Low	NT	May be declining at a moderately rapid rate, owing to bycatch on longline fisheries and perhaps the impacts of introduced predators. Threats and population status both remain poorly known.
Shy Albatross	<i>Thalassarche cauta</i>	3	Australia	A	14,618 (2005-2013)	↑	Low	NT	Breeds on just three islands and may be susceptible to stochastic events and human activities, although one nesting site is moderately widely separated from the other two.
White-capped Albatross	<i>Thalassarche steadi</i>	5	NZ	?	100,525 (1995-2013)	?	-	NT	The population trend of this albatross remains poorly known, as there is high inter-annual variability in breeding numbers and estimates prior to 2007 are not comparable with those made since. Analysis of recent data suggest that the trend may in fact be stable, but the species remains categorised as Near Threatened given the continuing uncertainty over its trend and because, given its longevity and slow productivity, and a high rate of mortality recorded in longline and trawl fisheries, it may be declining at a moderately rapid rate.
Northern giant Petrel	<i>Macronectes halli</i>	50		A	10,318 (1973-2013)	↑	Medium	LC	Had been predicted to undergo a moderately rapid population decline in the near future but has instead shown a significant increase during the past two decades (probably owing to greater availability of carrion from expanding populations of fur seals, increased waste from commercial fishing operations, and the use of measures to reduce seabird bycatch around some breeding colonies).
Southern giant Petrel	<i>Macronectes giganteus</i>	119		A	47,043 (1958-2013)	↑	Medium	LC	Recent analysis of trend data for the global population over the past three generations (64 years) gives a best case estimate of a 17 % increase and a worst case scenario of a 7.2 % decline (Chown <i>et al.</i> 2008 unpubl. report to SCAR).

<sup>1</sup> **Site:** usually an entire, distinct island or islet, or rarely, section of a large island (>3,000km<sup>2</sup>). ACAP database. <[data.acap.aq](http://data.acap.aq)>. May 2014.

<sup>2</sup> **Breeding Frequency:** A = Annual, B = Biennial

<sup>3</sup> **ACAP Trend:** ↑ increasing, ↓ declining, ↔ stable, ? unknown

<sup>4</sup> **IUCN Status:** CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern. IUCN 2013. IUCN Red List of Threatened Species. Version 2014.3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. 3 February 2015.