

Information of seabirds bycatch in area south of 25 S latitude in 2010 from 2015

Nobuhiro Katsumata<sup>1</sup>, Kotaro Yokawa<sup>1</sup> and Kazuhiro Oshima<sup>1</sup>

*SUMMARY*

*Seabird bycatch data collected by on-board observers in the area south of 25S in the Atlantic and the Indian Ocean in recent years is reviewed. Results revealed there is common tendency in between the southern bluefin tuna catch pattern and seabird bycatch pattern. Seabird bycatch pattern is also suggested to be influenced by geographical area as well as environmental condition. The results of this study is also indicate that the recent increasing trend of the nominal CPUE<sup>1</sup> of seabird is biased by the recent increase of the observer data in the area with higher seabird CPUE. Authors indicated these findings be considered in the future catch and effort data analysis.*

*KEYWORDS: Catch/effort, Longline, bycatch*

---

<sup>1</sup> National Research Institute of Far Seas Fisheries, Fisheries Research Agency, 5-7-1 Orido, Shimizu, Shizuoka, 424-8633, Japan.  
Email: katsumata@affrc.go.jp

## 1. Introduction

High latitudinal area of the south Atlantic and the south Indian Oceans are one of important regions for Japanese far seas longliners to catch high quality tunas like southern bluefin tuna. At the same time, longline operations in this region is known to have bycatch of seabirds, many of them need to be conserved due to their low stock condition. In the present study, observer reported catch and effort data of seabirds analyzed to the benefit of the evaluation of the impact of operations of Japanese longliners on seabird stocks.

## 2. Material and Methods

Same observer data used in the analysis of positive catch ratio of seabird (Inoue *et al.*, 2016). Those are mainly composed by CCSBT observers, and data collected by ICCAT and IOTC observers are also included. The data used in this analysis is limited to the one collect in the area south of 25S in the Atlantic and Indian Oceans where new seabird mitigation measures are mandated recently (ICCAT Rec11-09, IOTC Res12/06).

Species identification of bycaught seabirds were conducted by National Research Institute of Far Seas fisheries Japan and Birdlife International using photos taken by observers in the manner designated in the observer manual. For the simplification of the analysis, bycaught seabird species are classified into 13 groups shown in **Table 1**. In general, Japanese on-board observers to the longline boats cannot monitor whole gear retrieving process but only cover 80 – 90 % of it. Thus, the number of hooks whose gear retrieving is covered by observer in each set is used as the amount of effort for the analysis of catch and effort data of seabirds in this study. To analyze spatial pattern of bycatch, subareas are developed as shown in **Figure 1**, and quarter of the year is used for the analysis of seasonal pattern of bycatch.

Japanese observers are requested to report target species in each set through the questionnaire to the skipper, and this data is used for the target species determination. When multiple species are reported as target in single set, species having higher market value is adopted as the target species in this study. In the high latitudinal area of southern hemisphere in the Atlantic and Indian Oceans, southern bluefin tuna has highest market values for Japanese longliners, and second highest is bigeye tuna, 3<sup>rd</sup> is yellowfin tuna, 4<sup>th</sup> is albacore and values of other fishes are generally lower than these four tuna species.

## 3. Results and discussions

Almost sets monitored by on-board observers in the Atlantic was southern bluefin tunas in the period analyzed (**Table 2**). This is primarily due to the fact that almost observers dispatched to the Atlantic were CCSBT observers. Because some large number of other tunas than southern bluefin tuna are also caught in the south of 25S in the Atlantic (Yokawa *et al.*, 2016), this also indicates that sets targeted bigeye tuna, yellowfin tuna and albacore are not well monitored by observers. Sets targeting fishes other than southern bluefin tuna should also be monitored in the future. In the Indian Ocean, sets targeting other fishes than southern bluefin tuna seemed to be monitored to some extent (**Table 2**). The effect of target species on seabird bycatch is not sufficiently evaluated, but it should

be conducted because the result of **Table 2** indicates the fact that observers not randomly monitor sets from the view point of target fish.

The nominal CPUEs (n / 1000 hooks) showed general increasing trend in both Atlantic and Indian Oceans (**Figure 2**). The subarea specific nominal CPUE in two major fishing grounds of Japanese longliners in the south of 25S in the Atlantic and Indian Oceans (off South Africa and the southwest Indian Ocean) showed rather fluctuating patterns (**Figures 3 and 6**). One of the reason of this are the insufficient coverage of the data (**Figures 5 and 8**), which necessitate strengthening of monitoring, fluctuation of CPUEs are also seen in subareas some large number of data are continuously available (e.g. subareas 6 and 7 (ICCAT)). Investigation of the cause for the nominal CPUE observed in the subareas with relatively higher number of data would be important for the better understanding of the mechanism of seabird bycatch.

No clear and consistent increasing trends are not observed in the subarea specific CPUE trends (**Figures 3 and 6**), some notable high CPUEs can be observed in many subareas in both two major fishing ground, not only in most recent years but also in other years. At the same time in earlier years, some of the subareas had rather low or zero CPUE and this supposed to be due to insufficient number of data. One of the main reason of recent increase of the observer coverage should be due to the increase of allocation of southern bluefin tuna to Japan (CCSBT 2015). Seabird bycatch rate suggested to be higher in the higher latitudinal area (Inoue *et al.*, 2016) and southern bluefin tuna target set also conducted higher latitudinal area in the Southern Ocean (Yokawa *et al.*, 2016), the increase of quota of southern bluefin tuna should increase number of observer data in the area of higher seabird bycatch and would contributed to the increase of nominal CPUE. This indicates that the observed increasing trend of the nominal CPUEs in the Atlantic and Indian Oceans would be biased by the recent increase of observer data.

Considering the fact that the introduce of new mitigation major is believed to reduce seabird CPUE substantially (Melvin *et al.*, 2014), observed seabird CPUE level of Japanese longliners in most recent years is too high. National Research Institute of Far Seas Fisheries is now actively investigating its reason. This should be clarified as soon as possible to attain sound and prompt protection of seabirds.

In the off South Africa region, majority of seabird bycatch reported in the subareas south of 35S – 45S latitudinal band (**Figure 4**). In the Atlantic side (subareas 7 and 8 (ICCAT)), Grey-headed group and Black browed group were two major seabird groups, and wandering and Dark-colored groups also appeared constantly. In the Indian side, ratio of Grey-headed group decrease and White-chinned petrel appeared instead. Because catch species composition of Japanese longliners changes largely by relatively small geographical scale such as 5 degrees' latitude and this is suggested to be due to the influence of strong Agulhas warm current, relationship among seabird bycatch distribution pattern (Yokawa *et al.*, 2016), catch composition of longline, environmental condition and known seabird distribution pattern should be investigated to have better understanding of the mechanism of seabird bycatch occurrences. For this purpose, observer data before 2010 can also be used.

The bycatch composition of seabirds reported by Japanese observers drastically changed in the southwestern Indian Ocean (**Figure 7**). Yellow-nosed group is most popular in this region, and Gray-headed, Shy groups as

well as petrels seemed to appear constantly. In general, the level of CPUE of seabird seemed to be somewhat lower in this region from the one in the off South Africa (**Figures 3 and 6**), and this would be partially due to less variable environmental condition than one in the off South Africa region (Yokawa *et al.*, 2016).

Quarterly catch patterns by species group indicate that highest catch of seabird observed in 2<sup>nd</sup> quarter in subarea 6 (in the ICCAT region) where largest number of seabird recently caught in the off South Africa region (**Figure 9**). No apparent change of seasonal catch composition can be seen in between the period of 2010 – 2013 and 2014 – 2015, while the number of reported bycatch in the 3<sup>rd</sup> quarter increased in the later period. The largest catch of seabirds obtained in the 3<sup>rd</sup> quarter in subarea 6 and 14 (in the IOTC region) and some large number of bycatch also seen in the 2<sup>nd</sup> quarter (**Figures 10 and 11**). Catch species compositions are seemed not to be changed largely in between the period of 2010 – 2013 and 2014 – 2015, while the number of reported bycatch in the 2<sup>nd</sup> quarter increased in the later period.

In the present study, seabird bycatch data collected by on-board observers in the area south of 25S in the Atlantic and the Indian Ocean in recent years is quickly reviewed in the same strategy as the review of log-book data (Yokawa *et al.*, 2016). Results of this study revealed there is common tendency in between the southern bluefin tuna catch pattern and seabird bycatch pattern. Seabird bycatch pattern is also suggested to be influenced by geographical area as well as environmental condition. These finding would offer some important information for the catch and effort analysis of seabird bycatch, and thus, further study using additional data like the ones in before 2010 as well as ones of other fleets should be conducted.

## Reference

CCSBT. 2015. Report of the twenty second annual meeting of the commission. 12-15 October 2015, Yeosu, South Korea, 122pp.

Inoue, Y., M. Kanaiwa, M., Yokawa, K and Oshima, K. 2016. Examination of factors affecting seabird bycatch occurrence rate in southern hemisphere in Japanese longline fishery with using random forest. ICCAT SCRS/2016/162, 45p.

Inoue, Y., M. Kanaiwa, M., Yokawa, K., Okamoto, K and Oshima, K. 2016. Modeling of bycatch occurrence rate of seabirds for Japanese longline fishery operated in southern hemisphere. SCRS/2016/163, 13p.

Melvin E. F., Guy T. J., and Read L. B. 2014. Best practice seabird bycatch mitigation for pelagic longline fisheries targeting tuna and related species. Fisheries Research 149: 5-18.

Yokawa, K., Oshima, K., Inoue, Y and Katsumata, N. 2016. Operational pattern of Japanese longliners in the south of 25S in the Atlantic and the Indian Ocean for the consideration of seabird bycatches, ICCAT SCRS/2016/161, 14p.

**Table 1.** Grouping of seabirds used in this study.

Group name	Species name	Scientific name
Wandering group	Wandering Albatross	<i>Diomedea exulans</i>
	Tristan Albatross	<i>Diomedea dabbenena</i>
	Gibsoni Albatross	<i>Diomedea gibsoni</i>
	Antipodean Albatross	<i>Diomedea antipodensis</i>
Royal group	Southern Royal Albatross	<i>Diomedea epomophora</i>
	Northern Royal Albatross	<i>Diomedea sanfordi</i>
Black-browed group	Black-browed Albatross	<i>Thalassarche melanophris</i>
	Campbell Albatross	<i>Thalassarche impavida</i>
Yellow-nosed group	Atlantic Yellow-nosed Albatross	<i>Thalassarche chlororhynchos</i>
	Indian Yellow-nosed Albatross	<i>Thalassarche carteri</i>
Shy group	Shy Albatross	<i>Thalassarche cauta</i>
		<i>Thalassarche steadi</i>
	Salvin's Albatross	<i>Thalassarche salvini</i>
	Chatham Albatross	<i>Thalassarche eremita</i>
Grey-headed group	Grey-headed Albatross	<i>Thalassarche chrysostoma</i>
Buller's group	Buller's Albatross	<i>Thalassarche bulleri</i>
Dark-colored group	Sooty Albatross	<i>Phoebetria fusca</i>
	Light-mantled Albatross	<i>Phoebetria palpebrata</i>
Macronectes	Hall's Giant-Petrel	<i>Macronectes halli</i>
	Antarctic Giant-Petrel	<i>Macronectes giganteus</i>
White-chinned petrel		<i>Procellaria aequinoctialis</i>
Grey petrel		<i>Procellaria cinerea</i>
Flesh-footed shearwater		<i>Puffinus carneipes</i>
Other petrels		


**Table 2.** Ratio of sets by target species by year. Information reported by on-board observer was used. When multiple target species were reported.

ICCAT region


Species name	Scientific name	2010	2011	2012	2013	2014	2015
Southern bluefin tuna	<i>Thunnus maccoyii</i>	0	86.5	89.8	77.0	95.0	86.1
Bigeye tuna	<i>Thunnus obesus</i>	0	2.7	5.6	7.0	2.5	8.4
Yellowfin tuna	<i>Thunnus albacares</i>	0	0	0	3.5	0.3	0.3
Albacore	<i>Thunnus alalunga</i>	0	0	0	10.9	0.3	4.2
Other		100	10.8	4.6	1.6	1.9	1.1
Total operation		104	148	108	256	321	380

IOTC region

Species name	Scientific name	2010	2011	2012	2013	2014	2015
Southern bluefin tuna	<i>Thunnus maccoyii</i>	0	44.7	63.6	44.8	59.0	52.5
Bigeye tuna	<i>Thunnus obesus</i>	0	15.4	29.3	53.6	15.4	33.8
Yellowfin tuna	<i>Thunnus albacares</i>	0	0	1.7	0	1.2	9.8
Albacore	<i>Thunnus alalunga</i>	0	0	0	0.6	24.4	2.5
Other		100	39.8	5.4	1.1	0	1.3
Total operation		243	492	297	181	410	396




**Figure 1.** Subarea stratification in the Atlantic Ocean (upper) and Indian Ocean (lower) used in this study.




**Figure 2.** Nominal CPUE (n / 1000 hooks) of seabird by catch in two major fishing ground of Japanese longliners in the Atlantic and Indian Oceans. Left panel shows the CPUE in off South Africa (20W – 50E, 25S – 55S) and right panel shows one in the southwest Indian Ocean (70E – 120E).







**Figure 3.** Schematic figure of area specific nominal CPUE (n / 1000 hooks) trend of seabird in the period between 2010 – 2015 in off South Africa region. CPUE calculate by data collected by on-board observers for Japanese longliners.




**Figure 4.** Schematic figure of area specific catch number of seabird by species group in the period between 2010 – 2015 in off South Africa region. Data collected by on-board observers for Japanese longliners.




**Figure 5.** Schematic figure of area specific annual catch number of seabird (orange line) and annual number of hooks (blue bar) in the period between 2010 – 2015 in off South Africa region. Data collected by on-board observers for Japanese longliners.




**Figure 6.** Schematic figure of area specific catch number of seabird by species group in the period between 2010 – 2015 in the southwest Indian Ocean.. Data collected by on-board observers for Japanese longliners.




**Figure 7.** Schematic figure of area specific catch number of seabird by species group in the period between 2010 – 2015 in the southwest Indian Ocean. Data collected by on-board observers for Japanese longliners.




**Figure 8.** Schematic figure of area specific annual catch number of seabird (orange line) and annual number of hooks (blue bar) in the period between 2010 – 2015 in the southwest Indian Ocean. Data collected by on-board observers for Japanese longliners.



**Figure 9.** Quarterly bycatch number of seabird in subarea 6 in the ICCAT region (off South Africa) during the period of 2010 – 2013 (left) and 2014 – 2015 (right). Data collected by on-board observers was used.




**Figure 10.** Quarterly bycatch number of seabird in subarea 6 in the IOTC region (southwest Indian) during the period of 2010 – 2013 (left) and 2014 – 2015 (right). Data collected by on-board observers was used.



**Figure 11.** Quarterly bycatch number of seabird in subarea 14 in the IOTC region (southwest Indian) during the period of 2010 – 2013 (left) and 2014 – 2015 (right). Data collected by on-board observers was used.


APPENDIX; Annual Bycatch distribution map of seabird by species group in 2010 - 2015

ALZ: Large albatross, *Aptenodytes patagnicu*: king penguin.










Aves: Unidentified birds, DBN: Tristan albatross.




Latitude



Longitude





DIC: Grey-headed albatross, DIM: Black-browed albatross.



DIP: South royal albatross, DIX: Wandering albatross.



Latitude



Longitude

Macronectes spp: Unidentified giant petrel, MAH: Northern giant petrel.








Fig . continued. MAI: Southern giant petrel, PCI: Grey petrel.








PHE: Light-mantled albatross, PHU: Sooty albatross.



PRO: White-chinned petrel, Procellariidae: Unidentified petrel.




Latitude







Longitude

PUG: Great shearwater, Spheniscidae: Unidentified penguin.



Latitude



Longitude