CCSBT-ERS/2203/Annual Report – Japan (ERSWG Agenda Item 2.1)

National Report of Japan

Overview of Researches on Ecologically Related Species in Japanese SBT Longline Fishery, 2018-2020

Daisuke Ochi¹, Yasuko Semba¹, Yukiko Inoue¹, Tomoyuki Itoh¹, Sachiko Tsuji¹, Hiroshi Minami¹ and Masahiro Akiyama²

¹Fisheries Resources Institute, Japan Fisheries Research and Education Agency ²Fisheries Agency of Japan

要約

日本の漁業においてミナミマグロを対象とするのは、はえ縄漁業である。2018 年と2019 年、2020年におけるミナミマグロはえ縄漁業に従事した漁船数はそれぞれ、87、87、79 隻 であった。日本のミナミマグロはえ縄漁船が操業する水域は、CCSBT 統計海区の4、5、 7、8 及び9海区である。水産庁は、漁獲成績報告書の提出を船に義務付けると共に、1991年 からミナミマグロ漁獲量情報を収集するために漁業データ即時収集プログラム(RTMP)を実 施してきた。1995 年にはRTMPをすべてのミナミマグロはえ縄漁船を対象に実施してい る。

日本の科学オブザーバー計画は、1992年から開始されており、操業位置、漁獲努力量、 漁獲対象·非対象種の漁獲量、生物情報及び海鳥の偶発的捕獲などが本計画において記録さ れている。2018年と2019年、2020年におけるミナミマグロはえ縄漁船の科学オブザーバー 配乗隻数はそれぞれ、7隻と20隻、5隻であった。両年の科学オブザーバーによる調査カバ ー率はそれぞれ、隻数ベースでは8.1%と23.0%、6.4% で、総投下鈎数の6.4%と22.0%、 10.4%が観察された。本文書において、ミナミマグロはえ縄漁船に乗船した科学オブザーバ ーによって記録されたサメ類、海鳥類及び海亀類の捕獲数を報告した。

Summary

Japanese fleet is using only longline gear to catch southern bluefin tuna (SBT). Number of vessels engaging the SBT longline fishery was 87, 87, 79 in 2018, 2019 and 2020 respectively. Fishing grounds for SBT in recent years correspond to the CCSBT statistical areas of 4, 5, 7, 8 and to 9. Historically, logbook was submitted from fishermen to government as an obligation. In addition, Fisheries Agency of Japan started Real Monitoring Program (RTMP) from 1991 to monitor the catch of SBT. All the vessels for the SBT longline fishery have been monitored through this program since 1995.

Scientific observer program on the SBT fishery has been conducted by Japan since 1992, collecting information on fishing position, effort, catch of target and non-target species, biological information, incidental catch of seabirds, etc. The scientific observers were deployed to 7, 20 and 5 fishing vessels in 2018, 2019 and 2020, respectively. Coverage rates of observation were 8.1%, 23.0% and 6.4% for vessels and 6.4%, 22.0% and 10.4% for hooks in 2018, 2019 and 2020, respectively. This document reported captures of sharks, seabirds and sea turtles recorded by the scientific observers on-board the Japanese SBT longline vessels.

1. Introduction

Japanese fleet is using only longline gear to catch southern bluefin tuna (SBT). Since 1952, Japanese longline operation has started in the Indian Ocean that targeting yellowfin and bigeye tuna and caught, although SBT as was sub-target species for the longline fishery targeting yellowfin and bigeye tuna during the early stage of fishery. This is because of the fact that SBT in the tropical region were mostly spent of spawning with low meat quality so fishermen did not target it. Further south fishing grounds in the temperate waters for this species were developed in the late 1950s and 1960s. In addition, the innovation of super cold freezer has accelerated demand of "sashimi" grade SBT meat to the Japanese market. Recently the number of fishing vessels targeting SBT has gradually decreased due to the strong regulation for stock management and government policy to reduce number of longline vessels several times done in the past.

Regarding the incidental catch of seabirds, tori line was used voluntarily by the fishermen in the early 1990s, and the Government of Japan has introduced a mandatory measure for SBT longliners to use tori line since 1997. Research effort to modify tori line, weighted branchline and alternative methods possibly avoiding incidental catch of seabirds have continued. According to the international plans of action for reducing incidental catch of seabirds in longline fisheries and for the conservation and management of sharks, Japan established National Plans of Action in 2001 and has promoting mitigation of incidental take of seabirds, sea turtles and management of pelagic sharks.

2. Review of SBT Fisheries

Fleet size and distribution

The number of longline fishing vessels for SBT has been decreasing since the peak of about 300 in 1985. Fisheries Agency of Japan (FAJ) had reduced number of such vessels by 69 in 1981, 100 in 1982 and 132 in 1998. Vessel reduction policy in 1998 would have influenced further decline of number of vessels after then. The number of vessels has been less than 100 recently. Recent fishing grounds were off Cape of Good Hope (Area 9), southeastern Indian Ocean (Area 8), southeast of Australia (Area 4) and water near Tasmania Island (Area 7). Thus, the Japanese vessels were mainly operating in these areas, namely Area 4, 7, 8 and 9, in the second and third quarters for SBT.

Distribution of Catch and Effort

Catch and Effort data submitted to CCSBT were summarized. Effort of Japanese longline as the number of hooks used distributed widely in the southern hemisphere (Fig. 1). However, the major area of SBT catch came from Area 4, 7, 8 and 9 (Fig. 2).



Fig.1. Number of hooks of Japanese longline by 5x5 degrees square in 2018, 2019 and 2020.



Fig.2. Number of SBT caught by Japanese longline by 5x5 degrees square in 2018, 2019 and 2020.

3. Fisheries Monitoring for Each Fleet

Since 1991, FAJ has carried out Real Time Monitoring Program (RTMP) to monitor the catch of SBT. The number of vessels monitored by the program was 12-15 during 1991-1994, and all the vessels operating SBT fishing ground have been monitored by the RTMP since 1995. Each vessel sends daily reports including fishing position, effort, and catch by species in number and weight to the Fisheries Agency. The information is entered into the database in a short time.

Since 1992, Japan has conducted scientific observer program on SBT fishery and collected information including fishing position, effort, catch of target and non-target species, biological information, incidental catch of seabirds, etc. In 2018, 2019 and 2020, Japan deployed scientific observers to 7, 20 and 5 fishing vessels, respectively. While the observers were onboard the vessels, the vessels used 16,941, 13,277 thousand and 13,833 thousand hooks in 2018, 2019 and 2020, respectively. Coverage of observation was 8.1%, 23.0% and 6.4% for vessels and 6.4%, 22.0% and 10.4% for hooks in 2018, 2019 and 2020, respectively (Table 1).

Table 1. Number and coverage of cruises, sets and hooks observed in the Japanese RTMP observer program in 2018-2020. The number of hooks used by observed vessels in 2020 indicated the number of hooks observed in hauling operation and the numbers in brackets indicated the number of hooks used by observed vessels. Coverage for the number of hooks was calculated based on those numbers. This change was done since those coverages will be used for estimation of bycatch frequency from 2020.

Area	Calender year	Number of vessels observe d	Number of all vessels	Cover rate for the number of vessel	Number of hooks used by observed vessels (x 1000)	Number of hooks used by all vessels (x 1000)	Cover rate for the number of hook
Area 4	2018	1	22	4.5%	98	1,203	8.2%
	2019	5	24	20.8%	121	793	15.2%
	2020	0	15	0.0%	0	568	0.0%
Area 5	2018	0	7	0.0%	0	975	0.0%
	2019	1	7	14.3%	99	628	15.7%
	2020	1	9	11.1%	72 (94)	405	17.7% (23.1%)
Area 6	2020	1	4	25.0%	6 (7)	103	5.4% (7.1%)
Area 7	2018	1	27	3.7%	103	4,820	2.1%
	2019	6	26	23.1%	964	3,961	24.3%
	2020	1	21	4.8%	128 (168)	2,697	4.8% (6.2%)
Area 8	2018	3	21	14.3%	438	3,170	13.8%

	2019	4	21	19.0%	300	2,324	12.9%
	2020	1	31	3.2%	104 (116)	3,539	2.9% (3.3%)
Area	2018	3	39	7.7%	440	6,773	6.5%
9							
	2019	10	39	25.6%	1,438	5,571	25.8%
	2020	4	30	13.3%	721 (1,055)	6,521	11.1%
							(16.2%)
Total	2018	7	86	8.1%	1,079	16,941	6.4%
	2019	20	87	23.0%	2,921	13,277	22.0%
	2020	5	78	6.4%	1,031 (1,440)	13,833	7.5% (10.4%)

4. Seabird

Annual number of incidental catch of seabirds in the Japanese SBT longline fishery in 2018, 2019 and 2020 were updated based on the data collected through the scientific observer programs (see Table 2 and Appendix). Annual total captures were 10, 95 and 22 birds for large albatrosses, 21, 91 and 21 birds for dark colored albatrosses, 213, 997 and 67 birds for other albatrosses and, 7, 346 and 0 birds for unidentified albatrosses, 66, 455 and 62 birds for giant petrels, and 5, 7, and 10 birds for other seabirds, 1, 8 and 0 birds for unidentified seabirds in 2018, 2019 and 2020, respectively (Table 2).

5. Other Non-target Fish

The captures and mortalities of sharks in CCSBT fisheries are summarized in Table 2. 12, 14 and 10 species/species group of elasmobranchs were reported by the scientific observers in 2018, 2019 and 2020, respectively. Blue shark was dominant among elasmobranch catch observed, followed by porbeagle, and shortfin mako shark (CCSBT-ERS/2203/BGD 02, 03, 04).

Many teleosts were caught by longline fishery other than tunas and billfishes in the SBT fishing ground. There were 32, 49 and 32 species/species group of teleost fish including tuna and billfish found in the observer data in 2018, 2019 and 2020, respectively. Butterfly tuna, escoler, oilfish, opah, lancetfishes, sunfish and pomfrets were the major components of teleost catch (other than tuna and billfish) recorded in the in the high sea longline fishery (CCSBT-ERS/2203/BGD 02, 03, 04).

6. Marine Mammal and Marine Reptile

One capture of marine reptile in CCSBT fisheries was recorded in 2019 (Table 2). Five captures of marine mammals were recorded in 2019 (CCSBT-ERS/2203/BGD 02, 03, 04). Incidental catch of marine mammal and marine reptile occurred at a negligible level in the Japanese high-sea SBT longline fishery. There is not enough number of observations for the appropriate statistical estimation of the total incidental catch for these animals.

7. Mitigation Measures to Minimize Seabird and Other Species Bycatch

Current Measures

Mandatory measures

All tuna longline fishing vessels including those operating to catch SBT are

obliged to comply with respective rules adopted by the WCPFC, IATTC, IOTC and ICCAT, when operating in the Convention areas of these RFMOs. In addition, the Government of Japan has prepared law every time when there is amendment of the mitigation measures of these RFMOs and instructed to obey these measures for tuna longliners to obey these regulations.

Updated conservation and management measure to mitigate seabird bycatch were adopted at WCPFC and entered into force in December 2018 (WCPFC). Japan has amended its domestic regulations in compliance with currently active measures and implemented.

The measures that the Government of Japan to enforce and monitor the level of compliance for bycatch mitigation measures included a dispatch of enforcement vessels to the fishing areas, record of mitigation measures deployed through the logbook and collecting necessary information by scientific observers on board the operating vessels. The boarding observers and vessels carrying them are carefully selected so that avoiding the same vessels being selected in subsequent years. In addition to the mitigation measures adopted by each longline boat, Japanese observer program (JOP) has started to collect information of the general specifications of the mitigation measures adopted by each boat, such as the weight and position of swivels in the weighted branch line as well as the general configuration of tori lines, for the future detailed evaluation of the effect of mitigation measures.

Voluntary Measures, including information on proportion of fleet using the voluntary measures:

In February 2001, in accordance with "International Plan of Action for reducing incidental catch of seabirds in longline fisheries" of FAO, the Government of Japan developed "Japan's National Plan of Action for reducing incidental catch of seabirds in longline fisheries", in which FAJ instructed every fishermen to voluntarily carry out night setting, use of weighted branch line to ensure speedy precipitation of bait and use of properly defrozen bait in addition to the use tori lines which was already mandatory at that time.

Measures under Development/Testing

1) Mitigation measures:

Effectiveness of tori-line and weighted branchline by Japanese research vessel was examined in the North Pacific from April to May 2014-2016. The result indicated that tori-line and weighted branchlines are effective mitigation measures for tuna longline operations in the North Pacific. The further research on tori-line for small-scale vessels to reduce incidental catch of seabirds in the north Pacific has been carried out during 2017-2020 and use of light-weight materials for main line of tori line can be create enough aerial extent to prevent seabird attack to baited hooks.

2) Conservation and management

Large number of leatherback turtles is known to nest in Jamursba-medi and Wermon, West Papua, Indonesia. Nest counts, assessment of hatching success, and improvement of nesting environments for leatherbacks have been conducted since 1999 in Indonesia with the collaboration of the Indonesia Sea Turtle Research Center and Everlasting Nature of Asia, which is a Non-Profit Organization (NPO) in Japan. The nesting survey revealed that Indonesian population of leatherback turtles were suffering from poor reproductive success due to beach erosion, egg predation and low hatching rates. The Everlasting Nature constructed electric fences in the highest density nesting area to prevent pig predation on leatherback eggs. The electric fence drastically reduced the predation rates of eggs. Sea turtle populations have been affected by many factors on land and at sea (disappearance of nesting beaches, hatchling production, predation of eggs and turtles, interaction with fisheries such as trawl, gillnet, set-net, trap, purse-seine, and longline). Therefore, holistic management is necessary for the conservation of sea turtles, especially leatherback turtles.

8. Public Relations and Education Activities

Public Relation Activities

Educational materials, including booklets, pamphlets, video program (DVD/VHS), cartoons were prepared by FRA, the Global Guardian Trust (GGT), Japan NUS and the Organization for the Promotion of Responsible Tuna Fisheries (OPRT), and were distributed to fishermen and other parties related to fishing industry to explain theimportance of reducing incidental catch of seabirds and sea turtles.

- Identification guide for sharks, seabirds and sea turtles.
- Booklets and leaflets that illustrate methods for avoiding incidental catch and
- appropriate handling of seabirds and sea turtles;
- A guidebook which summarizes the NPOA-Seabirds and NPOA-Sharks.
- A video program (VHS and DVD) which explains mitigation measures to reduce longline interactions with seabirds and sea turtles.

Education

Japan Tuna Fisheries Cooperative Association has distributed weight on branch line to Japanese SBT longline vessels before the commencement of the fishing season.

Japan Tuna Fisheries Cooperative Association has held the workshops on seabird mitigation measures for captains, fish masters and owners of Japanese SBT longline vessels in collaboration with relevant organizations including the Birdlife International in Cape Town every year since 2019.

Japan Tuna Fisheries Cooperative Association will continue its efforts to further promote implementation of seabird mitigation measures by Japanese SBT longline vessels, including holding workshops, providing opportunities for fishers to be educated on seabird mitigation at Japanese port and distribution of seabird mitigation devices, subject to COVID-19.

<u>Observers</u>

Before the cruises, scientific observer candidates are obligated to take a training seminar. JOP held the training seminars twice a year to train scientific observers in usual year. During the training seminars, the candidates brushed up their knowledge and skills on research method, recording procedure and safety. Training included the practices of measuring the fish size and of collecting the biological samples. After the return from the commercial longline vessels, every observer reported their research activity. Their experiences and information have been used for the improvement of the observer program and next research activity. (CCSBT-ERS/2203/BGD 02, 03, 04).

9. Information on other ERS (non-bycatch) such as prey and predator species

Japan has been accumulating data of stomach contents of southern bluefin tuna in age-1 sampled in the SBT trolling monitoring survey (Itoh 2021).

10. Others

No other information.

11. Implementation of the NPOA-Seabirds and IPOA-Sharks

Japan developed its own National Plans of Action (NPOAs) for both seabirds and sharks in 2001 according to the FAO International Plans of Action (IPOAs) and revised them in 2016 taking into account the latest management measures taken by several RFMOs. FAJ disseminated the NPOAs to fishermen through local governments and fishermen's organizations. FAJ has reviewed implementation status of these two NPOAs and submitted its implementation reports to the FAO Committee on Fisheries (COFI) every two years since 2003.

Table 2: Reporting form for estimation of total mortality of ERS in CCSBT fisheries

Country	-		Japan		Y	ear (calen	ıdar year)	2	018	-								
		Total & (Observed Effo	ort ¹			Observed	Captures			Proportior	of observ	red effort	with speci	fic mitig	gation m	easures	
Stratum (CCSBT Statistical Areas or finer scale)	Human Observer / EM ²	Total Effort ³	Total Observed Effort ³	Observe r Coverag e^4	Species ⁵	Retained (dead)	Fate (<i>m</i> Discarded (dead)	(live)	Other ⁸	TP + NS ⁶	$TP + WB^6$	$NS + WB^6$	$TP + WB + NS^{6}$	TP ⁶	NS ⁶	WB ⁶	NIL	Others 7
4	OBS	1,203,461	98,140	8%	BSH	0	236	342	3	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	BSH													
6	OBS	0	0		BSH													
7	OBS	4,819,712	103,240	2%	BSH	0	242	315	6	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	BSH	1,180	491	638	800	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	BSH	375	303	394	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	SMA	0	5	10	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	SMA													

¹ Values in these shaded cells will be repeated for all species within a strata.

² Use codes OBS = Human observers, EM = Electronic monitoring. The ERSWG recognised that there was no agreement that EM replace the requirement for 10% observer coverage, and that the proposed inclusion of the option to report on EM results was not intended to imply any such agreement but only to clarify the source of any data that were reported.

³ For longline provide number of hooks, for purse seine provide number of sets.

⁴ For longline provide as a percentage of the number of hooks, for purse seine provide as a percentage of the number of shots.

⁵ Use FAO's 3 alpha species codes.

 $^{^{6}}$ TP = tori poles, NS = night setting, WB = weighted branchline.

⁷ Add extra columns for other categories of mitigation measures, if required.

⁸ All other captures not included in the columns for Retained (dead), Discarded (dead), and Released (live), e.g. released with undetermined life status.

6	OBS	0	0		SMA													
7	OBS	4,819,712	103,240	2%	SMA	0	1	1	1	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	SMA	12	0	1	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	SMA	3	0	0	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	POR	0	5	6	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	POR													
6	OBS	0	0		POR													
7	OBS	4,819,712	103,240	2%	POR	0	10	17	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	POR	0	145	84	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	POR	0	162	206	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	SHK	0	2	31	3	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	SHK													
6	OBS	0	0		SHK													
7	OBS	4,819,712	103,240	2%	SHK	0	2	23	2	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	SHK	0	12	9	2	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	SHK	0	53	313	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	DAL	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	

OBS 974.625 0 0% DAL OBS 0 0 DAL OBS 4,819,712 103,240 2% DAL 0% 0% 0 0 0 0 31% 0% 0% 0% 69% 0% 437,771 OBS 3,170,450 14% DAL 0 0 3% 0% 0% 0% 97% 0% 0% 0% 0 0 OBS 6,772,686 440,088 6% DAL 0 21 34% 0% 0% 66% 0% 0% 0 0 0% 0% OBS 1,203,461 98,140 8% LAL 0 1 0 0 32% 0% 68% 0% 0% 0% 0% 0% OBS 0% LAL 974,625 0 OBS 0 0 LAL OBS 4,819,712 103,240 2% LAL 0 0 2 0 31% 0% 0% 0% 69% 0% 0% 0% OBS 3,170,450 437,771 14% LAL 0 3 0 0 0% 0% 97% 0% 3% 0% 0% 0% OBS 6,772,686 440.088 6% LAL 0 3 1 0 34% 0% 0% 0% 66% 0% 0% 0% OBS 1,203,461 98,140 8% OAL 0 13 32% 0% 0% 0% 68% 1 0 0% 0% 0%

0

0

0

1

0

1

CCSBT-ERS/2203/Annual Report - Japan

0%

0%

0%

0%

0%

0%

31%

3%

34%

69%

97%

66%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0

0

0

13

18

166

-12-

OAL

OAL

OAL

OAL

OAL

5

6

7

8

9

4

5

6

7

8

9

4

5

6

7

8

9

OBS

OBS

OBS

OBS

OBS

974,625

0

4,819,712

3,170,450

6,772,686

0

0

103,240

437,771

440,088

0%

2%

14%

6%

4	OBS	1,203,461	98,140	8%	UAL	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	UAL													
6	OBS	0	0		UAL													
7	OBS	4,819,712	103,240	2%	UAL	0	0	0	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	UAL	0	2	0	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	UAL	0	4	1	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	OPT	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	OPT													
6	OBS	0	0		OPT													
7	OBS	4,819,712	103,240	2%	OPT	0	0	0	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	OPT	0	0	0	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	OPT	0	0	0	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	OSB	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	OSB													
6	OBS	0	0		OSB													
7	OBS	4,819,712	103,240	2%	OSB	0	0	0	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	OSB	0	0	0	0	3%	0%	0%	0%	97%	0%	0%	0%	

9	OBS	6,772,686	440,088	6%	OSB	0	5	0	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	USB	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	USB													
6	OBS	0	0		USB													
7	OBS	4,819,712	103,240	2%	USB	0	0	0	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	USB	0	1	0	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	USB	0	0	0	0	34%	0%	0%	0%	66%	0%	0%	0%	
4	OBS	1,203,461	98,140	8%	TTL	0	0	0	0	32%	0%	0%	0%	68%	0%	0%	0%	
5	OBS	974,625	0	0%	TTL													
6	OBS	0	0		TTL													
7	OBS	4,819,712	103,240	2%	TTL	0	0	0	0	31%	0%	0%	0%	69%	0%	0%	0%	
8	OBS	3,170,450	437,771	14%	TTL	0	0	0	0	3%	0%	0%	0%	97%	0%	0%	0%	
9	OBS	6,772,686	440,088	6%	TTL	0	0	0	0	34%	0%	0%	0%	66%	0%	0%	0%	
TOTAL		16,940,934	1,079,239	6%		1,570	1,919	2,397	817	21%	0%	0%	0%	79%	0%	0%	0%	

Table 2: Continued

C	Country	-		Japan		Y	'ear (caler	ndar year)	2	019	-								
			Total &	Observed Effo	ort ⁹			Observed	Captures			Proportior	of observ	ved effort	with speci	ific mitig	gation m	easures	
	Stratum (CCSBT Statistical Areas or finer scale)	Human Observer / EM ¹⁰	Total Effort ¹¹	Total Observed Effort ³	Observe r Coverag e ¹²	Species ¹³	Retained (dead)	Fate (<i>ni</i> Discarded (dead)	umbers) Released (live)	Other ¹⁶	TP + NS ¹⁴	$TP + WB^6$	NS + WB ⁶	$\begin{array}{c} TP \\ + WB \\ + NS^6 \end{array}$	TP ⁶	NS ⁶	WB ⁶	NIL	Others
	4	OBS	793,271	120,577	15%	BSH	23	22	254	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
	5	OBS	627,957	98,556	16%	BSH	37	0	20	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
	6	OBS	0	0	0%	BSH													
	7	OBS	3,961,222	963,709	24%	BSH	182	125	1,548	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
	8	OBS	2,324,044	299,681	13%	BSH	714	88	1,123	7	7%	19%	0%	1%	73%	0%	0%	0%	0%
	9	OBS	5,570,508	1,438,404	26%	BSH	2,612	387	2,055	8	10%	6%	0%	3%	74%	1%	0%	0%	2%
	4	OBS	793,271	120,577	15%	SMA	5	7	14	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
	5	OBS	627,957	98,556	16%	SMA	10	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%

⁹ Values in these shaded cells will be repeated for all species within a strata.

¹⁰ Use codes OBS = Human observers, EM = Electronic monitoring. The ERSWG recognised that there was no agreement that EM replace the requirement for 10% observer coverage, and that the proposed inclusion of the option to report on EM results was not intended to imply any such agreement but only to clarify the source of any data that were reported.

¹¹ For longline provide number of hooks, for purse seine provide number of sets.

¹² For longline provide as a percentage of the number of hooks, for purse seine provide as a percentage of the number of shots.

¹³ Use FAO's 3 alpha species codes. ¹⁴ TP = tori poles, NS = night setting, WB = weighted branchline.

¹⁵ Add extra columns for other categories of mitigation measures, if required.

¹⁶ All other captures not included in the columns for Retained (dead), Discarded (dead), and Released (live), e.g. released with undetermined life status.

6	OBS	0	0	0%	SMA													
7	OBS	3,961,222	963,709	24%	SMA	29	24	88	2	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	SMA	11	1	8	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	SMA	57	0	14	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	POR	0	1	9	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	POR	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	POR													
7	OBS	3,961,222	963,709	24%	POR	0	80	399	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	POR	0	113	493	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	POR	1	433	780	1	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	SHK	0	3	38	1	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	SHK	0	0	36	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	SHK													
7	OBS	3,961,222	963,709	24%	SHK	0	8	86	122	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	SHK	0	2	49	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	SHK	0	31	743	1	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	DAL	0	1	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%

5	OBS	627,957	98,556	16%	DAL	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	DAL													
7	OBS	3,961,222	963,709	24%	DAL	0	1	0	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	DAL	0	1	0	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	DAL	0	87	1	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	LAL	0	2	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	LAL	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	LAL													
7	OBS	3,961,222	963,709	24%	LAL	0	40	2	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	LAL	0	0	0	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	LAL	0	47	4	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	OAL	0	52	1	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	OAL	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	OAL													
7	OBS	3,961,222	963,709	24%	OAL	0	700	10	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	OAL	0	16	0	3	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	OAL	0	213	2	0	10%	6%	0%	3%	74%	1%	0%	0%	2%

4	OBS	793,271	120,577	15%	UAL	0	0	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	UAL	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	UAL													
7	OBS	3,961,222	963,709	24%	UAL	0	0	0	176	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	UAL	0	5	0	27	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	UAL	0	1	0	137	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	GPT	0	1	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	GPT	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	GPT													
7	OBS	3,961,222	963,709	24%	GPT	0	108	1	37	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	GPT	0	7	0	8	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	GPT	0	290	3	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	OSB	0	0	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	OSB	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	OSB													
7	OBS	3,961,222	963,709	24%	OSB	0	0	0	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	OSB	0	0	0	0	7%	19%	0%	1%	73%	0%	0%	0%	0%

9	OBS	5,570,508	1,438,404	26%	OSB	0	5	2	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	USB	0	0	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	USB	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	USB													
7	OBS	3,961,222	963,709	24%	USB	0	0	0	8	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	USB	0	0	0	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	USB	0	0	0	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
4	OBS	793,271	120,577	15%	TTL	0	0	0	0	26%	13%	0%	7%	54%	0%	0%	0%	0%
5	OBS	627,957	98,556	16%	TTL	0	0	0	0	0%	0%	8%	0%	0%	16%	25%	0%	50%
6	OBS	0	0	0%	TTL													
7	OBS	3,961,222	963,709	24%	TTL	0	0	1	0	19%	14%	0%	4%	63%	0%	0%	0%	0%
8	OBS	2,324,044	299,681	13%	TTL	0	0	0	0	7%	19%	0%	1%	73%	0%	0%	0%	0%
9	OBS	5,570,508	1,438,404	26%	TTL	0	0	0	0	10%	6%	0%	3%	74%	1%	0%	0%	2%
TOTAL		13,277,002	2,920,927	22%		3,681	2,902	7,784	538	13%	10%	0%	3%	67%	1%	1%	0%	3%

Table 2: Continued

C	Country	-]	Japan		Y	ear (caler	ıdar year)	2	020	-								
			Total & (Observed Effc	ort ¹⁷			Observed	Captures			Proportior	of observ	ved effort	with speci	fic miti	gation m	easures	
	Stratum (CCSBT Statistical Areas	Human Observer /	Total Effort ¹⁹	Total Observed	Observe r Coverag	Species ²¹	Retained	Fate (<i>ni</i> Discarded	umbers) Released	0.1 24	TP +	$TP + WB^6$	$NS + WB^6$	TP + WB	TP ⁶	NS ⁶	WB ⁶	NIL	Others 23
	or finer scale)	EM ¹⁰	ļ	Effort	e ²⁰		(dead)	(dead)	(live)	Other ²⁴	NS			+ NS°		<u> </u>			
	4	OBS	568,194	0	0%	BSH													
	5	OBS	404,818	71,600	18%	BSH	184	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
	6	OBS	103,320	5,605	5%	BSH	5	3	2	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
	7	OBS	2,697,048	128,246	5%	BSH	99	79	130	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
	8	OBS	3,538,838	104,080	3%	BSH	175	0	4	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
	9	OBS	6,520,563	721,504	11%	BSH	1,392	959	475	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
	4	OBS	568,194	0	0%	SMA													
	5	OBS	404,818	71,600	18%	SMA	0	1	4	1	0%	10%	17%	5%	0%	0%	68%	0%	0%

¹⁷ Values in these shaded cells will be repeated for all species within a strata.

¹⁸ Use codes OBS = Human observers, EM = Electronic monitoring. The ERSWG recognised that there was no agreement that EM replace the requirement for 10% observer coverage, and that the proposed inclusion of the option to report on EM results was not intended to imply any such agreement but only to clarify the source of any data that were reported.

¹⁹ For longline provide number of hooks, for purse seine provide number of sets.

²⁰ For longline provide as a percentage of the number of hooks, for purse seine provide as a percentage of the number of shots.

²¹ Use FAO's 3 alpha species codes. ²² TP = tori poles, NS = night setting, WB = weighted branchline.

²³ Add extra columns for other categories of mitigation measures, if required.

²⁴ All other captures not included in the columns for Retained (dead), Discarded (dead), and Released (live), e.g. released with undetermined life status.

6	OBS	103 320	5 605	5%	SMA	0	1	0	0	0%	10%	0%	00%	0%	0%	0%	0%	0%
0	005	105,520	5,005	570	SMA	0	1	0	0	0 70	1070	0 70	9070	070	070	070	0 70	070
7	OBS	2,697,048	128,246	5%	SMA	0	15	3	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	SMA	0	0	1	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	SMA	0	7	9	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	POR													
5	OBS	404,818	71,600	18%	POR	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	POR	0	1	5	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	POR	0	27	123	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	POR	0	0	22	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	POR	0	550	537	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	SHK													
5	OBS	404,818	71,600	18%	SHK	0	1	30	8	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	SHK	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	SHK	0	1	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	SHK	0	0	2	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	SHK	0	51	181	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	DAL													

5	OBS	404,818	71,600	18%	DAL	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	DAL	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	DAL	0	0	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	DAL	0	0	0	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	DAL	0	21	0	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	LAL													
5	OBS	404,818	71,600	18%	LAL	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	LAL	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	LAL	0	1	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	LAL	0	0	0	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	LAL	0	14	7	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	OAL													
5	OBS	404,818	71,600	18%	OAL	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	OAL	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	OAL	0	6	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	OAL	0	0	0	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	OAL	0	56	5	0	43%	18%	0%	6%	32%	0%	0%	0%	0%

4 OBS 568.194 0 0% UAL 5 OBS 404,818 71,600 18% UAL 0 0 0 0 0% 10% 17% 5% 0% 0% 68% 0% 0% 103,320 5,605 UAL 0 0% 6 OBS 5% 0 0 0 0% 10% 0% 90% 0% 0% 0% 0% 7 OBS 2,697,048 128,246 UAL 0 0 31% 0% 0% 0% 0% 5% 0 0 0% 69% 0% 0% 8 OBS 3,538,838 104,080 3% UAL 0 0 24% 6% 0% 68% 0% 0 0 0% 0% 2% 0% 9 OBS 6,520,563 721,504 UAL 0 0 0 0 43% 18% 0% 32% 11% 6% 0% 0% 0% 0% OBS 0% GPT 4 568,194 0 5 OBS 404,818 71,600 18% GPT 0 0 0 0 68% 0% 10% 17% 5% 0% 0% 0% 0% 103,320 5,605 5% GPT 0 0 0 0 10% 0% 90% 0% 6 OBS 0% 0% 0% 0% 0% 7 OBS 2,697,048 128,246 5% GPT 0 6 0 0 0% 69% 0% 0% 31% 0% 0% 0% 0% 8 OBS 3,538,838 104,080 3% GPT 0 0 0 0 24% 6% 0% 2% 68% 0% 0% 0% 0% OBS 6,520,563 721,504 11% GPT 52 18% 0% 32% 9 0 4 0 43% 6% 0% 0% 0% 0% 4 OBS 568,194 0 0% OSB 5 OBS 404,818 71,600 18% OSB 0 0 0 0 0% 10% 17% 5% 0% 0% 68% 0% 0% 103,320 OSB 10% 0% 0% 0% 6 5,605 0 0 0 0 OBS 5% 0% 90% 0% 0% 0% 7 OBS 2,697,048 128,246 OSB 0 0 0 0 31% 0% 69% 0% 0% 5% 0% 0% 0% 0% OBS 3,538,838 104,080 OSB 68% 8 3% 0 0 0 0 24% 6% 0% 2% 0% 0% 0% 0%

9	OBS	6,520,563	721,504	11%	OSB	0	6	4	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	USB													
5	OBS	404,818	71,600	18%	USB	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	USB	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	USB	0	0	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	USB	0	0	0	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	USB	0	0	0	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
4	OBS	568,194	0	0%	TTL													
5	OBS	404,818	71,600	18%	TTL	0	0	0	0	0%	10%	17%	5%	0%	0%	68%	0%	0%
6	OBS	103,320	5,605	5%	TTL	0	0	0	0	0%	10%	0%	90%	0%	0%	0%	0%	0%
7	OBS	2,697,048	128,246	5%	TTL	0	0	0	0	0%	31%	0%	69%	0%	0%	0%	0%	0%
8	OBS	3,538,838	104,080	3%	TTL	0	0	0	0	24%	6%	0%	2%	68%	0%	0%	0%	0%
9	OBS	6,520,563	721,504	11%	TTL	0	0	0	0	43%	18%	0%	6%	32%	0%	0%	0%	0%
TOTAL		13,832,781	1,031,035	7%		1,855	1,858	1,548	9	33%	18%	1%	14%	29%	0%	5%	0%	0%

References

Itoh, T. (2021) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2020/2021. CCSBT-ESC/2108/29.

Appendix

Group names	English names	2018	2019	2020
Large albatrosses	Wandering albatross	8	62	20
	Tristan albatross	0	1	2
	Gibson's albatross	1	7	0
	Wandering albatross group	1	22	0
	Southern royal albatross	0	1	0
	Royal albatross group	0	2	0
Dark colored albatrosses	Sooty albatross	15	59	18
	Light-mantled albatross	5	32	3
	Unidentified dark colored albatrosses	1	0	0
Other albatrosses	Black-browed albatross	51	57	19
	Campbell albatross	4	53	0
	Atlantic yellow-nosed albatross	0	0	1
	Indian yellow-nosed albatross	2	22	0
	Black-browed albatross group	14	111	5
	Buller's albatross group	14	339	2
	Shy-type albatrosses	13	342	11
	Yellow-nosed albatross group	0	1	0
	Grey-headed albatross	114	64	29
	Unidentified other albatrosses	1	8	0
Unidentified albatrosses		7	346	0
Giant petrels	Northern giant petrel	3	31	11
	Southern giant petrel	7	27	6
	Southern fulmar	0	1	0
	Grey petrel	29	35	11
	White-chinned petrel	6	169	4
	Parkinson's petrel	0	2	6
	Spectacled petrel	0	2	0
	Flesh-footed shearwater	1	4	0
	Great shearwater	11	136	24
	Sooty shearwater	1	0	0
	Unidentified giant petrels	2	1	0
	Unidentified petrels	6	47	0
Other seabirds		5	7	10
Unidentified seabirds		1	8	0