# INDONESIAN SCIENTIFIC OBSERVER PROGRAM ACTIVITIES IN INDIAN OCEAN FROM 2015-2017

Bram Setyadji<sup>1</sup>, Zulkarnaen Fahmi<sup>1</sup>, Toni Ruchimat<sup>2</sup>

<sup>1)</sup> Research Institute for Tuna Fisheries, Bali, Indonesia <sup>2)</sup> Research Institute for Marine Fisheries, Jakarta, Indonesia

Prepared for the Extended Scientific Committee for the Twenty Third Meeting of the Scientific Committee, San Sebastian, Spain, 3 - 8 September, 2018

#### Summary

This paper presents a progress summary of the Indonesian scientific observer program on the tuna fishing vessels operating in the Indian Ocean. The observer data is the most detailed information not only associated with catch and effort, but also information on fishing practices, gear configuration and environmental conditions. Only low fleet coverages were available from this data set. Hence this could higher the possibility to get robust abundance indices from the fishery.

#### Indonesian observer program in the Indian Ocean

There are two Indonesian observer programs currently operating in the Indian Ocean, namely national observer program and scientific observer program. The national observer program has been formally legalized by the MMAF since 2013 through Ministerial Decree No. 1/2013. Catch and effort data have been collected through the national observer program, however, issues on data entry and validity as well as the need for verification and validation prior analysis is remained (Irianto et al. 2016). The need to improve the capacity of the observers and related officers were highlighted (Irianto et al. 2016).

The scientific observer program has initiated since 2005 through a collaboration between RCCF-MMAF (Indonesia) and CSIRO (Australia) funded by ACIAR on a trial observer program. This program was then continued and funded by the Indonesian Government through the RCCF between 2009 and 2011. The Research Institute for Tuna Fisheries (under RCCF) was established in 2011 in order to conduct researches on tuna fisheries in the Indian Ocean including continuing the scientific observer program. The trial observer program (2005-2008) has been reported in the

13<sup>th</sup> Meeting of the Extended Scientific Committee in 2008, while the scientific observer data have been reported within the Indonesian National Report Scientific Meetings of the IOTC and CCSBT since 2010. This information paper provides an update of the Indonesian scientific observer program activities in the Indian Ocean in 2016.

### Design of the scientific observer program

There are 8 trained scientific observers at the RITF. The observers were not only deployed in the longline vessels operating out of Benoa, but also Muara Baru and Cilacap. In 2017, there were four scientific observers' trips in total, with total day-at-sea ranged from 49 to 69 fishing days per trip with total efforts of 192,188 hooks and the number of sets ranged between 22 and 37 sets. The observer coverages were 3.6% in terms of total active vessels. Geographically, the capacity of scientific observer covered the fishing ground of statistical area 1 and 2.

Table 1.Observer activities in authorized Fishing Vessels in 2017 (source: Indonesian National<br/>Report for the CCSBT SC meeting)

Trip	Base	Date	DAS	No of Settings	No of Hooks	No of SBT	HR (x10 <sup>3</sup> )	CCSBT Statistical Area
1	Cilacap	16/09/17	69	22	48,112	0	0.000	1
2	Cilacap	16/09/17	63	37	42,180	0	0.000	1
3	Cilacap	18/10/17	51	29	37,936	0	0.000	1
4	Benoa	10/11/17	49	32	63,960	197	3.080	2

### Spatial distribution of the observed sets and tuna catch composition

The observed tuna longline sets covered the area between  $0.651^{\circ}$ -33.676° S and 75.788°-129.522° E and concentrated in the area above 20° S (area 1) and below 25° S (area 2). Darker color indicates higher number of hooks observed (Figure 1). Southern bluefin tunas were caught in set within region CCSBT statistical area 2 (Figure 2)




Figure 1. Map of the distribution of Indonesian tuna longline efforts year 2017, acquired from scientific observer data.




Figure 2. Spatial distribution of tuna catch composition in 2017, acquired from scientific observer data.

### Trends of catch and effort recorded by the observers

Over the last five years, the effort observed by the scientific observers are declining. Mainly because technical issues related to some implementation on onboard observer's regulations recently. In 2017, only 4 trips were deployed (Table 2). Observed catch and effort from scientific observer data shown the hook rates for most tuna species were declining in the last 5 years (Figure 3.). In order to get a robust catch and effort data, Indonesia is currently improving the quality and quantity of logbook data.

SBT									
TRIP	n	HR	Length (cmTL)						
			Average	Min	Max				
1	0	0.000	-	-	-				
2	0	0.000	-	-	-				
3	0	0.000	-	-	-				
4	197	3.080	157.58	135	197				

Table 2. Estimated hook-rates of SBT on authorized fishing vessels




Figure 3. Nominal CPUE of longline catch from 2005-2017 based on scientific observer data

## Challenges

Main challenge in term of improvement of the use of the scientific observer data are related to the following issues: low spatial and temporal coverages (region), low fleet coverage (gear types) and no replication for each observed vessel. The scarce dataset was cause by reluctance from tuna longline fishing company to be monitored by scientific observer, they still worry, some of the importance information, such as fishing ground would be exposed to the public.

Year	No. Of Obs	No. Of Trips	No. Of Company	Total Day at Sea	Days/Trip	Avg (d/trip)
2005	6	6	1	251	19-22	20
2006	6	19	5	758	7-99	39
2007	6	14	5	648	21-108	34
2008	5	15	7	481	23-66	30
2009	5	14	8	535	15-59	38
2010	5	8	4	240	40-50	50
2011	5	6	3	210	30-50	40
2012	6	7	5	496	33-208	83
2013	5	3	3	170	52-60	57
2014	8	6	4	371	29-90	62
2015	4	5	5	241	31-61	48
2016	3	3	3	170	32-86	57
2017	4	4	2	232	49-69	58

Appendix 1. Annual activities of scientific observer based in Benoa Bali from 2005 to 2017

Appendix 2.	Observed catch and effort data of ecologically related species (ERS) from scientific
	observer program in 2016.

Trip	Total Hooks	MON	BSH	BTH	CCE	CDF	DAV	DCU	DOL	EIL	FAL
1	640,620	-	0.186	-	-	0.197	0.813	-	-	-	-
2	1,090,830	0.114	2.005	-	-	0.128	-	0.015	-	-	-
3	1,088,500	0.087	0.600	0.059	-	0.167	1.358	-	0.021	-	0.022
4	898,304	0.051	0.685	-	0.025	0.203	1.593	-	-	0.025	0.025
Trip	Total Hooks	GES	ISB	LEC	LKV	LOP	MSO	NGA	OCS	OIL	PDM
1	640,620	-	0.304	1.182	0.051	-	-	4.662	-	0.068	-
2	1,090,830	-	-	0.230	-	-	-	1.113	-	0.186	0.265
3	1,088,500	0.245	0.463	2.456	0.059	0.022	0.087	6.880	0.022	0.043	-
4	898,304	0.355	0.076	1.578	-	-	-	5.428	0.076	0.203	-
Trip	Total Hooks	PHU	PSK	PTH	SMA	TCR	TPA	TSK	TST	WAH	
1	640,620	-	0.203	-	-	0.467	-	-	1.166	0.175	
2	1,090,830	0.015	0.374	0.030	0.559	-	-	-	0.102	-	
3	1,088,500	-	0.815	-	-	0.248	0.059	-	0.538	0.326	
4	898,304	-	0.434	-	-	0.254	0.101	0.152	0.621	0.180	

Code	Species	2013	2014	2015	2016	2017
ALZ	Albatrosses nei	0	0	7	0	0
BAR	Barracudas nei	0	4	5	6	0
BSH	Blue shark	37	63	137	105	184
BTH	Bigeye thresher					3
BTS	Hound needlefish	1	0	0	1	0
BRZ	Pomfrets, ocean breams nei	194	232	121	0	0
CCB	Spinner shark	4	17	1	3	0
CCE	Bull shark					1
CCL	Blacktip shark	0	0	1	0	0
ССР	Sandbar shark	0	0	0	0	0
DAV	Sind danio	914	511	206	272	157
DKK	Leatherback turtle	0	0	0	0	0
DOL	Common dolphinfish	11	15	7	13	33
EBS	Brilliant pomfret	5	0	0	0	1
FAL	Silky shark	0	0	26	0	2
GES	Snake mackerel					26
HAR	Longnose chimaeras	22	3	14	46	0
ISB	Cookie cutter shark					39
LEC	Escolar	284	666	490	353	240
LHX	Seagulls nei	0	1	0	0	0
LKV	Olive ridley turtle	6	12	1	15	5
LMA	Longfin mako	0	0	0	0	0
LAG	Opah	51	29	30	60	13
MOX	Ocean sunfish	11	3	2	1	0
MSK	Mackerel sharks, porbeagles nei	1	0	0	0	0
ALX	Long snouted lancetfish	1738	921	739	693	796
OCS	Oceanic whitetip shark	2	8	4	4	4
OIL	Oilfish	7	58	16	8	24
PHU	Sooty albatross	0	0	0	0	1
PSK	Crocodile shark	48	90	108	174	84
PTH	Pelagic thresher	0	0	0	0	2
RME	Longhorned mobula	0	1	1	0	0
RMJ	Spinetail mobula	1	1	0	0	0
SMA	Shortfin mako	2	2	1	5	35
SPL	Scalloped hammerhead	0	0	1	0	0

**Appendix 3.** List of ecologically related species incidentally caught by Indonesia tuna longliners in the last 5 years (2013-2017)

Code	Species	2013	2014	2015	2016	2017
SPY	Hammerhead sharks, etc. nei	1	0	0	0	0
SSH	Scarlet shrimp	0	1	0	0	0
TCR	Keeltail pomfret	91	90	45	62	42
TIG	Tiger shark	1	0	0	0	0
TPA	Dealfish	0	0	0	0	7
TSK	Flapnose houndshark	1	2	6	1	6
TST	Sickle pomfret	60	110	29	117	105
TTH	Hawksbill turtle	0	0	0	0	0
TTX	Marine turtles nei	0	0	0	0	0
TUG	Green turtle	0	0	0	0	0
WAH	Wahoo	60	96	63	61	30