


This paper presents the outcomes of a review of seabird bycatch mitigation measures for pelagic longline fisheries conducted by ACAP's Seabird Bycatch Working Group (SBWG) at its meeting in Guayaquil, Ecuador, held from 29 August – 2 September 2011. The review is presented in two parts. The first consists of a summary of the review, providing concise advice on the efficacy of the mitigation measures reviewed. The second part contains detailed information on the review conducted on each of the mitigation measures.

PART 1



Goal: Reduce the bycatch of seabirds to the lowest possible level.

Summary

Recognising that most (84%) breeding albatrosses overlap with the pelagic longline fisheries for tuna and swordfish managed by the five tuna Regional Fisheries Management Organisations (RFMOs), the adoption of best practice seabird conservation in these fisheries is a high priority for the Agreement on the Conservation of Albatrosses and Petrels (ACAP) (AC3 Info 18, 2007).

A combination of weighted branchlines, bird scaring lines and night setting are best practice mitigation in pelagic longline fisheries. These measures should be applied in high risk areas

such as the high latitudes of southern hemisphere oceans and lower to mid-latitude fisheries of both the northern and south east Pacific to reduce the incidental mortality to the lowest possible levels. Other factors such as safety, practicality and the characteristics of the fishery should also be recognised.

Currently, no single mitigation measure can reliably prevent the incidental mortality of seabirds in most pelagic longline fisheries. The most effective approach is to use the above measures in combination.

Introduction

The incidental mortality of seabirds, mostly albatrosses and petrels, in longline fisheries continues to be a serious global concern and was major reason for the establishment of the Agreement on the Conservation of Albatrosses and Petrels (ACAP). In longline fisheries seabirds are killed when they become hooked and drowned while foraging for baits on longline hooks as the gear is deployed. They also can become hooked as the gear is hauled; however, many of these seabirds can be released alive with careful handling. Although most mitigation measures are broadly applicable, the application and specifications of some will vary with local longlining methods and gear configurations. For example, most scientific literature on seabird bycatch mitigation in pelagic fisheries relates to larger vessels, with little research attention to smaller vessels and the gear configuration and methods of artisanal fleets; seabird bycatch mitigation advice is under development. ACAP has comprehensively reviewed the scientific literature dealing with seabird bycatch mitigation in pelagic fisheries and this document is a distillation of that review (AC6 Final Report ANNEX 13).

BEST PRACTICE MEASURES

1. Branchline weighting

Branchlines should be weighted to sink the baited hooks rapidly out of the diving range of feeding seabirds. Weighted lines sink faster and more consistently, resulting in dramatic reductions in seabird attacks on baited hooks and seabird mortality; no negative effect has been demonstrated on the catch rate of fishes. Continued refinement of line weighting configurations (mass, number and position of weights and materials) through controlled research and application in fisheries, is encouraged to find configurations that are most safe, practical and effective.

Scientific studies have demonstrated that branchline weighting configurations with more mass close to the hook, sinks the hooks most rapidly and consequently is most effective at reducing seabird interactions and mortalities. Current recommended minimum standard for branchline weighting configurations are the following:

Greater than 45 g attached within 1 m of the hook or;

Greater than 60 g attached within 3.5 m of the hook or;

Greater than 98 g weight attached within 4 m of the hook.

Positioning weight farther than 4 m from the hook is not recommended.

2. Night setting

Setting longlines at night, between nautical twilight and nautical dawn, is highly effective at reducing incidental mortality of seabirds because the majority of vulnerable seabirds are inactive at night.

3. Bird scaring lines

Properly designed and deployed bird scaring lines deter birds from sinking baits, thus dramatically reducing seabird attacks and related mortalities. A bird scaring line is a line that runs from a high point at the stern to a device or mechanism that creates drag at its terminus. As the vessel moves forward, drag lifts the section of line closest to the vessel from the water into the air. Brightly coloured streamers hanging from the aerial extent of the line scare birds from flying to and under the line preventing them from reaching the baited hooks. It is the aerial extent (out of water) section with suspended streamers that scares birds from the sinking baits.

Bird scaring lines should be the lightest practical strong fine line. Lines should be attached to the vessel with a barrel swivel to minimise rotation of the line from torque created as it is dragged behind the vessel.

Towed objects, applied to increase drag, and with it bird scaring line aerial extent, are prone to tangling with float lines leading to lost bird scaring lines, interruptions in vessel operations and in some cases lost fishing gear. Alternatives, such as adding short streamers to the in-water portion of the line, can enhance drag while minimising tangles with float lines. Weak links (breakaways) should be incorporated into the in-water portion of the line safety and operational problems should lines become tangled.

Given operational differences in pelagic longline fisheries due to vessel size and gear type, bird scaring lines specifications have been divided into recommendations for vessels greater than 35 metres and those less than 35 metres.

3. (a) Recommendations for vessels >35 m total length

Simultaneous use of two bird scaring lines, one on each side of the sinking longline, provide maximum protection from bird attacks under a variety of wind conditions and are recommended as best practice for larger vessels.

Bird scaring lines should include the following specifications:

Bird scaring lines should be deployed to maximise the aerial extent. Aerial extent is a function of vessel speed, height of the attachment point to the vessel, drag, and weight of bird scaring line materials.

Vessels should deploy bird scaring lines with a minimum aerial extent of 100 m.

Streamers should be: brightly coloured, a mix of long and short streamers, placed at intervals of no more than 5 m, and long streamers attached to the line with swivels that prevent streamers from wrapping around the line. All streamers should reach the sea-surface in calm conditions.

Baited hooks shall be deployed within the area bounded by the two bird scaring lines. Bait-

casting machines shall be adjusted so as to land baited hooks within the area bounded by the bird scaring lines.

If large vessels use only one bird scaring line, the bird scaring line should be deployed windward of sinking baits. If baited hooks are set outboard of the wake, the bird scaring line attachment point to the vessel shall be positioned several meters outboard of the side of the vessel that baits are deployed. This position is best achieved using a purpose build davit (tori pole) located as close to the stern and as far aft as practical. Proper outboard positioning also minimises the likelihood of bird scaring lines tangling on float lines.

3. (b) Recommendations for vessels <35 m total length

A single bird-scaring line using either long and short streamers, or short streamers only, has been found effective on smaller vessels.

Streamers should be brightly coloured. Short streamers (>1 m) should be placed at 1 m intervals along the length of the aerial extent. Two designs have been shown to be effective: a mixed design that includes long streamers placed at 5 m intervals over the first 55 m of the bird scaring line and a design that does not include long streamers.

Vessels should deploy bird scaring lines with a minimum aerial extent 75 m.

Other Considerations

Area and seasonal closures:The temporary closure of important foraging areas (e.g. areas adjacent to important seabird colonies during the breeding season when large numbers of aggressively feeding seabirds are present) to fishing will eliminate incidental mortality of seabirds in that area.

Mainline tension:Setting mainline, branch lines and baited hooks into propeller turbulence (wake) slows sink rates and should be avoided.

Live vs. dead bait: Use of live bait should be avoided. Individual live baits can remain near the water surface for extended periods (e.g. up to 120 seconds), thus increasing the likelihood of seabird captures.

Bait hooking position:Baits hooked in either the head (fish), or tail (fish and squid), sink significantly faster than baits hooked in the mid-back or upper mantle (squid).

Offal and discard discharge management: Seabirds are attracted to discards, offal and used baits. Used baits should be retained during line hauling. Ideally offal and used baits should be discharged on the side of the vessel opposite of line hauling. Offal and discards should not be discharged during line setting. All hooks should be removed and retained on board before discards are discharged from the vessel.

New Technologies

New technologies that set or release baited hooks at depth (underwater setting device) or disarm hooks to specific depths, which have the potential to prevent seabird access to baits, are currently under development and undergoing sea trials.

Mitigation Technologies that are Not Recommended

Line shooters: There is no experimental evidence that line shooters reduce seabird bycatch in pelagic longline fisheries; therefore, they should not be considered a seabird bycatch mitigation option.

Olfactory deterrents: Olfactory deterrents (fish oils) have not been demonstrated to prevent or reduce seabird mortalities in pelagic longline fisheries.


Hook size and design: Changes to hook size and design may reduce the chance of seabird mortality in longline fisheries, but have not been sufficiently researched.

Side setting: Although side setting (defined as setting station a minimum of one metre forward of the stern and in combination with branchline weighting and a bird curtain) is being used in the Hawaiian surface longline fishery, it has not been tested in other fisheries, including southern hemisphere fisheries, consequently it cannot be recommended at this time.

Blue dyed bait: Blue dyed squid bait has been insufficiently researched and cannot be recommended.

Bait thaw status: In practical terms the thaw status of baits has no effect on the sink rate of baited hooks set on weighted lines.

PART 2

 <p>Agreement on the Conservation of Albatrosses and Petrels</p>	<h3>ACAP REVIEW OF SEABIRD BYCATCH MITIGATION MEASURES FOR PELAGIC LONGLINE FISHERIES</h3> <p><i>Amended at the Sixth Meeting of the Advisory Committee Guayaquil, Ecuador, 29 August – 2 September 2011</i></p>
---	--

Weighted branchlines, bird scaring streamer lines and night setting are best practice seabird bycatch mitigation in pelagic longline fisheries. ACAP's Seabird Bycatch Working Group (SBWG) has comprehensively reviewed the scientific literature dealing with seabird bycatch mitigation in pelagic fisheries and this document is a distillation of that review.

BEST PRACTICE MEASURES	
1.	Branchline weighting
2.	Night setting
3.	a). Bird scaring streamer lines for vessels > 35m in total length
3.	b). Bird scaring streamer lines for vessels <35m in total length
OTHER CONSIDERATIONS	
4.	Side setting with line weighting and bird curtain
5.	Blue dyed bait
6.	Line shooter
7.	Bait caster
8.	Underwater setting chute
9.	Management of offal discharge
10.	Live bait
11.	Bait thaw status
12.	Area closures

BEST PRACTICE MEASURES

1. Branchline weighting

Scientific evidence for effectiveness in pelagic fisheries

PROVEN AND RECOMMENDED. Should be used in combination with night setting and bird scaring lines. Brothers 1991; Boggs 2001; Sakai *et al.* 2001; Brothers *et al.* 2001; Anderson & McArdle 2002; Gilman *et al.* 2003a, Hu *et al.* 2005.

Caveats /Notes

Weights will shorten but not eliminate the zone behind the vessel in which birds can be caught. Even in demersal fisheries where weights are much heavier, weights must be combined with other mitigation measures (e.g. CCAMLR Conservation Measure 25-02).

Need for combination

Should be combined with bird scaring lines and night setting

Research needs

Mass and position of weight both affect sink rate. Further research on the effect of a range of weighting regimes on seabird mortality and catch rates of target and non-target fishes is needed (as has been completed for demersal [Spanish system] fisheries). Continued work to identify branchline weighting configurations (mass, placement, shape, number of leads, and materials) that are effective at reducing seabird bycatch with and without other mitigation, and that are safe and practical. Effect of propeller turbulence on baited hook sink rate and seabird mortality need to be quantified.

Minimum standards

Current minimum standards for branchline weighting configurations are:

Greater than 45 g attached within 1 m of the hook or;

Greater than 60 g attached within 3.5 m of the hook or;

Greater than 98 g weight attached within 4 m of the hook.

Positioning weight farther than 4 m from the hook is not recommended.

These regimes have been adopted in the Hawaiian (45 g at 1 m) and Australian (60 g at 3.5 m and 98 g at 4 m) pelagic longline fisheries and latter two regimes have been adopted by the Western and Central Pacific Fishing Commission (the WCPFC provisions also include the option of branchlines being configured with weights of 45 g to 60 g within 1 m of the hook). NB. The 98 g weights specified in the Australian fishery pertain to the line weighting experiment of Robertson *et al.* 2010. The commercially available leaded swivels used in the experiment weighed 98 g (not 100 g).

Implementation monitoring

Coastal state fisheries (vessels <35 m total length): Line weights crimped into branch lines technically very difficult to remove at sea. Inspection before departure from port of all gear bins on vessels considered an acceptable form of implementation monitoring.

Distant water fisheries (vessels >35 m total length): Technically possible to remove and/or re-configure gear at sea. Implementation monitoring by monitoring line sets using appropriate methods (e.g., observer inspection of line setting operations; video surveillance; at-sea compliance checks). Video surveillance conditional on mainline setter being fitted with motion sensors to trigger cameras.

2. Night setting

Scientific evidence for effectiveness in pelagic fisheries

PROVEN AND RECOMMENDED. Should be used in combination with weighted branch lines and bird scaring lines. Duckworth 1995; Brothers *et al.* 1999; Gales *et al.* 1998; Klaer&Polacheck 1998; Brothers *et al.* 1999; McNamara *et al.* 1999; Gilman *et al.* 2005; Baker & Wise 2005; Jiménez *et al.* 2009.

Caveats /Notes

Less effective during full moon, under intensive deck lighting or in high latitude fisheries in summer. Less effective on nocturnal foragers e.g. White-chinned Petrels (Brothers *et al.* 1999; Cherelet *et al.* 1996).

Need for combination

Should be used in combination with bird scaring lines and weighted branch lines

Research needs

Determine effectiveness of bird scaring lines and branchline weighting at night by characterising seabird behaviour at night using thermal or night vision technologies.

Minimum standards

Night defined as between nautical twilight and nautical dawn.

Implementation monitoring

Requires VMS (satellite transmitter) or fishery observers. Vessel speed and direction vary between transiting, line setting, line hauling and when vessels are stationary on fishing grounds. VMS-derived assessment of vessel activity in relation to time of nautical dawn and dusk considered acceptable for implementation monitoring. Alternatively VMS-linked sensors fitted to mainline setting and hauling drum could be used to indicate compliance, as could sensors to trigger video surveillance cameras. This facility is currently unavailable and requires development.

3	a). Bird scaring streamer lines for vessels > 35m in total length
----------	---

Scientific evidence for effectiveness in pelagic fisheries

PROVEN AND RECOMMENDED. For vessels > 35 m in length two streamer lines is considered best practice. Streamer lines with the appropriate aerial extent can be more easily rigged on large vessels. Two streamer lines are considered to provide better protection of baited hooks in crosswinds (Melvin *et al.* 2004; Melvin *et al.* 2011). Hybrid tori lines (with long and short streamers) were more effective than short tori lines (only short streamers) in deterring diving seabirds (white-chinned petrels) (Melvin *e.al.* 2010; Melvin *et al.* 2011).

Caveats /Notes

Potentially increased likelihood of entanglement, particularly if attachment points on davits (tori poles) are insufficiently outboard of vessels. Development of a towed device to prevent tangling with fishing gear essential to improve adoption and compliance.

Diving species increase vulnerability of surface foragers (albatrosses) due to secondary interactions.

Need for combination

Should be used with appropriate line weighting and night setting.

Research needs

Compare the effectiveness of one versus two bird scaring lines, including with respect to both primary and secondary interactions; develop methods that create drag to maximise aerial extent while minimising entanglements of the in-water portion of bird scaring lines with longline floats; and compare the effectiveness of bird scaring lines with different streamer lengths, configurations, and materials.

Minimum standards

Vessels should deploy bird scaring lines with a minimum aerial extent of 100 m. Streamers should be: brightly coloured, a mix of long and short streamers, placed at intervals of no more than 5 m, and long streamers attached to the line with swivels that prevent streamers from wrapping around the line. All streamers should reach the sea-surface in calm conditions.

If large vessels use only one streamer line it should be set to windward of sinking baits. If baited hooks are set outboard of the wake, the streamer line attachment point to the vessel should be positioned several meters outboard of the side of the vessel that baits are deployed.

Baited hooks shall be deployed within the area bounded by the two streamer lines. Bait-casting machines shall be adjusted so as to land baited hooks within the area bounded by streamer lines

Implementation monitoring

Requires fisheries observers, video surveillance, or at-sea surveillance (e.g. patrol boats or

aerial over-flights).

3	b). Bird scaring streamer lines for vessels <35m in total length
----------	--

Scientific evidence for effectiveness in pelagic fisheries

PROVEN AND RECOMMENDED. Imber 1994; Uozomi& Takeuchi 1998; Brothers *et al.* 1999; Klaer&Polacheck 1998; McNamara *et al.* 1999; Boggs 2001; CCAMLR 2002; Minami &Kiyota 2004; Melvin 2003. For vessels < 35 m in length a single BSL in combination with night setting and appropriate line weighting has been found effective for mixed and short streamer bird-scaring lines (ATF 2011; Domingo *et al.*, Gianucaet *al.* 2011).

Caveats /Notes

Development of a towed device to prevent tangling with fishing gear essential to improve adoption and compliance.

Diving species increase vulnerability of surface foragers (albatrosses) due to secondary interactions.

Need for combination

Should be used with appropriate line weighting and night setting.

Minimum standards

Vessels should deploy bird scaring lines with a minimum aerial extent 75 m. Streamers should be brightly coloured. Short streamers (>1 m) should be placed at 1 m intervals along the length of the aerial extent. Two designs have been shown to be effective: a mixed design that includes long streamers placed at 5 m intervals over the first 55 m of the bird scaring line and a design that does not include long streamers. Bird scaring lines should be the lightest practical strong fine line. Lines should be attached to the vessel with a barrel swivel to minimise rotation of the line from torque created as it is dragged behind the vessel.

Towed devices to create drag can tangle with float lines leading to interruptions in vessel operations and in some cases lost fishing gear. Short streamers can be tied into the line to bristle the line and create a bottlebrush like configuration to generate drag while minimising the chance of fouling streamer lines on float lines. Breakaways should be incorporated into the streamer line in-water extent to minimise safety and operational problems should a longline float foul or tangle with the in-water extent of a streamer line.

Implementation monitoring

Requires fisheries observers, video surveillance, or at-sea surveillance (e.g. patrol boats or aerial over-flights).

OTHER CONSIDERATIONS

4. Side setting with line weighting and bird curtain

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN AND NOT RECOMMENDED FOR SOUTHERN HEMISPHERE FISHERIES. Brothers & Gilman 2006; Yokota & Kiyota 2006.

Caveats /Notes

Only effective if hooks are sufficiently below the surface by the time they reach the stern of the vessel and protected by a bird curtain. In Hawaii, side-setting trials were conducted with bird curtain and 45-60 g weighted swivels placed within 0.5 m of hooks. Japanese research concludes must be used with other measures (Yokota & Kiyota 2006). Not tested in southern hemisphere fisheries and cannot be recommended at this time.

Need for combination

Lines set from the side of vessels must be appropriately weighted and protected by an effective bird curtain. Requires thorough testing in southern hemisphere fisheries.

Research needs

Currently untested in southern hemisphere fisheries against assemblages of diving seabirds (e.g. *Procellaria* sp. Petrels and *Puffinus* sp. Shearwaters) and albatrosses - urgent need for research.

Minimum standards

Clear definition of side setting is required. As noted, side setting trials in Hawaii were conducted in conjunction with a bird curtain and 45-60 g leaded swivel < 1 m of the baited hook. Hawaiian definition is a minimum of only 1 m forward of the stern, which is likely to reduce effectiveness. The distance forward of the stern refers to the position from which baits are manually deployed. Baited hooks must be thrown by hand forward of the bait deployment location if they are to be afforded "protection" by being close to the side of the vessel.

Implementation monitoring

Requires fisheries observers or video surveillance.

5. Blue dyed bait

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN AND NOT RECOMMENDED. Boggs 2001; Brothers 1991; Gilman *et al.* 2003a; Minami & Kiyota 2001; Minami & Kiyota 2004; Lydon & Starr 2005. Cocking *et al.* 2008.

Caveats /Notes

New data suggests only effective with squid bait (Cocking *et al.* 2008). Onboard dyeing requires labour and is difficult under stormy conditions. Results inconsistent across studies.

Need for combination

Must be combined with bird scaring lines or night setting.

Research needs

Need for tests in Southern Ocean.

Minimum standards

Mix to standardised colour placard or specify (e.g. use 'Brilliant Blue' food dye (Colour Index 42090, also known as Food Additive number E133) mixed at 0.5% for minimum 20 minutes).

Implementation monitoring

The current practice of dyeing bait on board vessels at sea requires observer presence or video surveillance to monitor implementation. Assessment of implementation in the absence of on-board observers or video surveillance requires baits be dyed on land and monitored through port inspection of all bait on vessels prior to departure on fishing trips.

6. Line shooter

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN AND NOT RECOMMENDED. Robertson *et al.* 2010.

Caveats /Notes

Mainline set into propeller turbulence with a line shooter without tension astern (e.g. slack) as in deep setting significantly slows the sink rates of hooks (Robertson *et al.* 2010). Use of a line shooter to set gear deep cannot be considered a mitigation measure.

Need for combination

Not Applicable.

Research needs

Not Applicable.

Minimum standards

Use of this measure is not recommended as a mitigation measure.

Implementation monitoring

Not Applicable.

7. Bait caster

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN AND NOT RECOMMENDED. Duckworth 1995; Klaer&Polacheck 1998.

Caveats /Notes

Not a mitigation measure unless casting machines are available with the capability to control the distance at which baits are cast. This is necessary to allow accurate delivery of baits under a bird scaring line. Current machines (without variable power control) likely to deploy baited hooks well beyond the streaming position of streamer lines, increasing risks to seabirds. Few commercially-available machines have variable power control. Needs more development.

Need for combination

Not recommended as a mitigation measure at this time.

Research needs

Develop (and implement) casting machine with a variable power control.

Minimum standards

Not recommended as a mitigation measure

Implementation monitoring

Not Applicable

8. Underwater setting chute

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN AND NOT RECOMMENDED. Brothers 1991; Boggs 2001; Gilman *et al.* 2003a; Gilman *et al.* 2003b; Sakai *et al.* 2004; Lawrence *et al.* 2006.

Caveats /Notes

For pelagic fisheries, existing equipment not yet sturdy enough for large vessels in rough seas. Problems with malfunctions and performance inconsistent (e.g. Gilman *et al.* 2003a and Australian trials cited in Baker & Wise 2005).

Need for combination

Not recommended for general application at this time.

Research needs

Design problems to overcome.

Minimum standards

Not yet established

Implementation monitoring

Not Applicable.

9. Management of offal discharge

Scientific evidence for effectiveness in pelagic fisheries

UNPROVEN. McNamara *et al.* 1999; Cherelet *et al.* 1996.

Caveats /Notes

Supplementary measure. Definition essential. Offal attracts birds to vessels and where practical should be eliminated or restricted to discharge when not setting or hauling. Strategic discharge during line setting can increase interactions and should be discouraged. Offal retention and/or incineration may be impractical on small vessels.

Need for combination

Must be combined with other measures.

Research needs

Further information needed on opportunities and constraints in pelagic fisheries (long and short term).

Minimum standards

Not yet established for pelagic fisheries. In CCAMLR demersal fisheries, discharge of offal is prohibited during line setting. During line hauling, storage of waste is encouraged, and if discharged must be discharged on the opposite side of the vessel to the hauling bay.

Implementation monitoring

Requires offal discharge practices and events to be monitored by fisheries observers or video surveillance.

10. Live bait

Scientific evidence for effectiveness in pelagic fisheries

LIVE BAIT NOT RECOMMENDED. Trebilcoet *et al.* 2010; Robertson *et al.* 2010.

Caveats /Notes

Live fish bait sinks significantly slower than dead bait (fish and squid), increasing the exposure of baits to seabirds. Use of live bait is associated with higher seabird bycatch rates.

Need for combination

Use of live bait is not a mitigation measure.

Research needs

Not Applicable.

Minimum standards

Live bait is not a mitigation measure.

Implementation monitoring

Not Applicable.

11. Bait thaw status

Scientific evidence for effectiveness in pelagic fisheries

NOT RECOMMENDED. Brothers 1991; Duckworth 1995; Klaer&Polacheck; Brothers *et al.*1999; Robertson & van den Hoff 2010.

Caveats /Notes

Baits cannot be separated from others in frozen blocks of bait, and hooks cannot be inserted in baits, unless baits are partially thawed (it is not practical for fishers to use fully frozen baits). Partially thawed baits sink at similar rates to fully thawed baits.

Need for combination

Not a mitigation measure

Research needs

Not Applicable.

Minimum standards

Not recommended as a mitigation measure.

Implementation monitoring

Not Applicable.

12. Area closures

Scientific evidence for effectiveness in pelagic fisheries

PROVEN AND RECOMMENDED. Avoiding fishing at peak areas and during periods of intense foraging activity has been used effectively to reduce bycatch in longline fisheries.

Caveats /Notes

An important and effective management response, especially for high risk areas, and when other measures prove ineffective. Highly effective for target locations/seasons but may displace fishing effort into adjacent or other areas which may not be as well regulated, thus leading to increased incidental mortality elsewhere.

Need for combination

Must be combined with other measures, both in the specific areas when the fishing season is opened, and also in adjacent areas to ensure displacement of fishing effort does not merely lead to a spatial shift in the incidental mortality.

Research needs

Further information about the seasonal variability in patterns of species abundance around fisheries.

Minimum standards

No work done but highly recommended.

Implementation monitoring

Vessels equipped with VMS and activities monitored by appropriate management authority is considered appropriate monitoring. Areas/seasons should be patrolled to ensure effectiveness if IUU activities are suspected.

REFERENCES

- Anderson, S. and McArdle, B., 2002. Sink rate of baited hooks during deployment of a pelagic longline from a New Zealand fishing vessel. *New Zealand Journal of Marine and Freshwater Research*, 36: 185–195.
- ATF 2011. Developments in experimental mitigation research – Pelagic longline fisheries in Brazil, South Africa and Uruguay. Sixth Meeting of Advisory Committee. *Guayaquil, Ecuador, 29 August – 2 September 2011*.
- Baker, G.B., and Wise, B.S. 2005. The impact of pelagic longline fishing on the flesh-footed shearwater *Puffinus carneipes* in Eastern Australia. *Biological Conservation*, 126: 306–316.
- Boggs, C.H., 2001. Deterring albatrosses from contacting baits during swordfish longline sets. In: Melvin, E., Parrish, J.K. (Eds), *Seabird Bycatch: Trends, Roadblocks and Solutions*. University of Alaska Sea Grant, Fairbanks, Alaska, pp. 79–94.
- Brothers, N. and Gilman, E. 2006. Technical assistance for Hawaii-based pelagic longline vessels to modify deck design and fishing practices to side set. Prepared for the National Marine Fisheries Service Pacific Islands Regional Office. Blue Ocean Institute, September 2006.

- Brothers, N.P. 1991. Approaches to reducing albatross mortality and associated bait loss in the Japanese long-line fishery. *Biological Conservation*, 55: 255–268.
- Brothers, N., Gales, R. and Reid, T. 1999. The influence of environmental variables and mitigation measures on seabird catch rates in the Japanese tuna longline fishery within the Australian Fishing Zone 1991-1995. *Biological Conservation*, 88: 85–101.
- Brothers, N., Gales, R., and Reid, T., 2001. The effect of line weighting on the sink rate of pelagic tuna longline hooks, and its potential for minimising seabird mortalities. CCSBT-ERS/0111/53.
- Brouwer, S. and Walker, N. 2008. Use of light streamer lines and line weighting on longline vessels and the implications for seabird bycatch. WCPFC Scientific Committee Fourth Regular Session, 11-22 August 2008 WCPFC-SC4-2008/EB-IP-3.
- CCAMLR, 2002. Report of the working group on fish stock assessment. Report of the twenty-first meeting of the Scientific Committee of the Commission for the Conservation of Marine Living Resources. Commission for the Conservation of Marine Living Resources, Hobart.
- Cherel, Y., Weimerskirch, H. and Duhamel., G 1996. Interactions between longline vessels and seabirds in Kerguelen Waters and a method to reduce seabird mortality. *Biological Conservation*, 75: 63–70.
- Cocking, L.J., Double, M.C., Milburn, P.J. and Brando, V.E. 2008. Seabird bycatch mitigation and blue-dyed bait: A spectral and experimental assessment. *Biological Conservation*, 14: 1354–1364.
- Dimas, G., Peppes, F., César, J., Marques, C., and Neves, T. 2011. The effect of leaded swivel position and light toriline on bird attack rates in Brazilian pelagic longline. Sixth Meeting of Advisory Committee. *Guayaquil, Ecuador, 29 August – 2 September 2011*.
- Domingo, A., Jiménez, S., Abreu, M., Forselledo, R., and Pons, M. 2011. Effectiveness of tori-line use to reduce seabird bycatch in the Uruguayan pelagic longline fleet. Sixth Meeting of Advisory Committee. *Guayaquil, Ecuador, 29 August – 2 September 2011*.
- Duckworth, K., 1995. Analysis of factors which influence seabird bycatch in the Japanese southern bluefin tuna longline fishery in New Zealand waters, 1989–1993. New Zealand Fisheries Assessment Research Document 95/26.
- Gales, R., Brothers, N. and Reid, T. 1998. Seabird mortality in the Japanese tuna longline fishery around Australia, 1988-1995. *Biological Conservation*, 86: 37–56.
- Gilman, E., Brothers, N., Kobayashi, D. R., Martin, S., Cook, J., Ray, J., Ching, G., and Woods, B. 2003a. Performance assessment of underwater setting chutes, side setting, and blue-dyed bait to minimise seabird mortality in Hawaii longline tuna and swordfish fisheries. Final report. Western Pacific Regional Fishery Management Council. Honolulu, Hawaii, USA. 42pp.
- Gilman, E., Boggs, C. and Brothers, N. 2003b. Performance assessment of an underwater setting chute to mitigate seabird bycatch in the Hawaii pelagic longline tuna fishery. *Ocean and Coastal Management*, 46: 985–1010.

- Gilman, E., Brothers, N. and Kobayashi, D. 2005. Principles and approaches to abate seabird bycatch in longline fisheries. *Fish and Fisheries*, 6: 35–49.
- Hu, F., Shiga, M., Yokota, K., Shiode, D., Tokai, T., Sakai, H., and Arimoto, T. 2005. Effects of specifications of branch line on sinking characteristics of hooks in Japanese tuna longline. *Nippon Suisan Gakkaishi* 71: 33–38.
- Imber, M.J., 1994. Report on a tuna long-lining fishing voyage aboard Southern Venture to observe seabird by-catch problems. Science & Research Series 65. Department of Conservation, Wellington, New Zealand.
- Jiménez S, Domingo A, and Brazeiro A. 2009. Seabird bycatch in the Southwest Atlantic: interaction with the Uruguayan pelagic longline fishery. *Polar Biology*, 32: 187–196.
- Klaer, N. and Polacheck, T. 1998. The influence of environmental factors and mitigation measures on by-catch rates of seabirds by Japanese longline fishing vessels in the Australian region. *Emu*, 98: 305–16.
- Lawrence, E., Wise, B., Bromhead, D., Hindmarsh, S., Barry, S., Bensley, N. and Findlay, J. 2006. Analyses of AFMA seabird mitigation trials – 2001 to 2004. Bureau of Rural Sciences. Canberra.
- Lokkeborg, S., 2003. Review and evaluation of three mitigation measures - bird-scaring line, underwater setting and line shooter - to reduce seabird bycatch in the north Atlantic longline fishery. *Fisheries Research*, 60: 11–16.
- Lydon, G. and Starr, P., 2005. Effect of blue dyed bait on incidental seabird mortalities and fish catch rates on a commercial longliner fishing off East Cape, New Zealand. Unpublished Conservation Services Programme Report, Department of Conservation, New Zealand. 12p.
- McNamara B, Torre L, and Kaaialii G. Hawaii longline seabird mortality mitigation project. Honolulu, HI, USA: Western Pacific Regional Fishery Management Council, 1999.
- Melvin, E. F., Guy, T. J. and Reid, L. B. 2010. Shrink and Defend: A Comparison of Two Streamer Line designs in the 2009 South Africa Tuna Fishery. Third Meeting of the Seabird Bycatch Working Group, ACAP, SBWG-3 Doc 13.rev1.
- Melvin, E. F., Sullivan, B., Robertson, G. and Wienecke, B. 2004. A review of the effectiveness of streamer lines as a seabird bycatch mitigation technique in longline fisheries and CCAMLR streamer line requirements. *CCAMLR Science*, 11: 189–201.
- Melvin, E.F. 2003. Streamer lines to reduce seabird bycatch in longline fisheries. Washington Sea Grant Program, WSG-AS 00-33.
- Melvin, E.F., Parrish, J.K., Dietrich, K.S. and Hamel, O.S. 2001. Solutions to seabird bycatch in Alaska's demersal longline fisheries. Project A/FP-7, WSG-AS 01-01, Washington Sea Grant.
- Minami, H. and Kiyota, M. 2001. Effect of blue-dyed bait on reducing incidental take of seabirds. *CCSBT-ERS/0111/61*. 7pp.
- Minami, H. and Kiyota, M., 2004. Effect of blue-dyed bait and tori-pole streamer on reduction of incidental take of seabirds in the Japanese southern bluefin tuna longline fisheries. *CCSBT-ERS/0402/08*.

- Robertson, G., Candy, S.G. and Wienecke, B. 2010. Effect of line shooter and mainline tension on the sink rates of pelagic longlines and implications for seabird interactions. *Aquatic Conservation: Marine and Freshwater Ecosystems* DOI: 10.1002/aqc.1100.
- Robertson, G., and van den Hoff, J. 2010. Static water trials on the sink rates of baited hooks to improve understanding of sink rates estimated at sea. Report to the Third meeting of the Seabird Bycatch Working Group of ACAP.
- Robertson, G., Candy, S. G., Wienecke, B., and Lawton, K. submitted, 2010. Experimental determinations of factors affecting the sink rates of baited hooks to minimise seabird mortality in pelagic longline fisheries.
- Sakai, H., Fuxiang, H., and Arimoto, T., 2004. Underwater setting device for preventing incidental catches of seabirds in tuna longline fishing, CCSBT-ERS/0402/Info06.
- Sakai, H., Hu, F., and Arimoto, T. 2001. Basic study on prevention of incidental catch of seabirds in tuna longline. CCSBT-ERS/0111/62.
- Trebilco, R., Gales, R., Lawrence, E., Alderman, R., Robertson, G. and Baker, G.B. 2010 (in press). Seabird bycatch in the Eastern Australian Tuna and Billfish pelagic longline fishery: temporal, spatial and biological influences. *Aquatic Conservation: Marine and Freshwater Ecosystems*.
- Uozumi, Y. and Takeuchi, Y. 1998. Influence of tori pole on incidental catch rate of seabirds by Japanese southern bluefin tuna longline fishery in high seas. CCSBT-WRS/9806/9 revised. 5pp.
- Yokota, K. and Kiyota, M. 2006. Preliminary report of side-setting experiments in a large sized longline vessel. WCPFC-SC2-2006/EB WP-15. Paper submitted to the Second meeting of the WCPFC Ecosystem and Bycatch SWG. Manila, 10th August 2006
- Yokota, K., Minami, H. and Kiyota, M. 2008. Direct comparison of seabird avoidance effect between two types of tori-lines in experimental longline operations. WCPFC Scientific Committee Fourth Regular Session, 11-22 August 2008 WCPFC-SC4-2008/EB-WP-7.